

spline functions were used to obtain the numerical solutions of the Fredholm integral equations of the second kind. In paper, [13], the wavelet-Galerkin method for the numerical solution of the Fredholm linear integral equations and the second-order integro-differential equations are discussed. A construction of a quadratic spline-wavelet basis on the unit interval, such that the wavelets have three vanishing moments and the shortest support among such wavelets was proposed, [13]. In paper, [14], a new collocation technique for the numerical solution of the Fredholm, Volterra, and mixed Volterra-Fredholm integral equations of the second kind is introduced, and a numerical integration formula on the basis of the linear Legendre multi-wavelets is also developed. The linear Legendre multi-wavelets basis for the proposed method is used. In this technique, the unknown function is approximated by the truncated linear Legendre multi-wavelets series, [14].

Good results are obtained by using the Chebyshev polynomials. Paper, [15], focused on fuzzy Fredholm integral equations of the second kind. Using the Chebyshev polynomials due to their smoothness and reasonable behavior near boundaries, a new method is proposed to solve the fuzzy Fredholm integral equation. In paper, [16], the approximate solution of linear Fredholm integral equations of the second type on a closed interval is studied. The Galerkin method enhanced with the Chebyshev polynomials was used to improve the approximate solution.

We also note the following papers. In paper, [17], the authors have used the advanced multistep and hybrid methods to solve the Volterra integral equation. In paper, [18], the forward-jumping methods of the hybrid type are used for the construction of the methods with a high order of accuracy. In paper, [19], the Half-Sweep Gauss-Seidel iteration, which was used to find the approximate solution of the fuzzy Fredholm integral equations of the second kind, was applied. In paper, [20], linear Volterra-Fredholm integral equations of the second kind were considered in reproducing kernel space. A new scheme with a high convergence order for solving the approximate solutions to oscillation and non-oscillation of exact solutions was proposed.

In paper, [21], a new technique is offered to solve three types of linear integral equations of the 2nd kind, including the Volterra-Fredholm integral equations (as a general case), the Volterra integral equations, and the Fredholm integral equations (as special cases). The new technique depends on

approximating the solution to a polynomial of degree $(m - 1)$ is described.

This work is a continuation of a series of works on the use of continuous local polynomial splines for solving interpolation problems and for solving integral equations, [22], [23]. This paper is devoted to the investigation of the stability of the solution of the integral equation using these local splines. As is known, the solution of integral equations of the second kind is reduced to finding the frame of the approximate solution. This means that we find approximations to the values of the function at the nodes of the grid (grid function). Usually, the integral equation is replaced by some difference scheme with a given order of accuracy. The approximate values of the function is converged to the values of the function at the grid nodes if the two conditions are fulfilled. These conditions are as follows: an approximation of the equation with a difference scheme, and the difference scheme is stable. Let for an approximate solution of the integral equation be

$$Au = f$$

a difference scheme is constructed

$$A_h u^h = f^h.$$

This scheme approximates the original equation with some order of accuracy. Suppose a linear normed space of functions defined on the grid is considered. The operator A_h maps the space U_h to the space F_h . A difference scheme is said to be stable on the right-hand side if, for any $f^h \in F_h$, the equation

$$A_h u^h = f^h$$

has a unique solution $u^h \in U_h$ and

$$\| u^h \|_{U_h} \leq C \| f^h \|_{F_h}.$$

Here C is a constant. Next, consider the definition

$$\| u^h \|_{U_h} = \max_i |u^h(x_i)|.$$

Thus, the convergence of the approximate solution to the values of the function at the grid nodes follows from the approximation of the original equation and the stability of the solution.

In the works of, [24], conditions for the stability of the solution of the Fredholm integral equation of

the second kind using the trapezoidal method were obtained. Thus, the convergence of the obtained approximate solution of the Fredholm integral equation of the second kind to the values of the function at the grid nodes was proved.

Next, we construct a numerical scheme to solve the Fredholm integral equation of the second kind.

This paper discusses the stability of the solution when we use the local splines of the second and seventh order of approximation. We use these splines if the kernel and the right side are sufficiently smooth functions. To construct an approximate solution at the points between the grid nodes, we use the interpolation of the same local splines or the integral equation with an obtained solution at the nodes.

2 Problem Formulation

Let $\{x_j\}$ be a grid of ordered nodes on the interval $[a, b]$: $a = x_0 < \dots < x_n = b$. Note that the approximations with the splines are constructed separately for each grid interval $[x_j, x_{j+1}]$. Let us assume that the values of the function $u(x)$ are given at the grid nodes. The approximation using basis splines is built separately on each grid interval as the sum of the products of the values of the function u at the grid nodes and the basis splines ω_j .

Let r, r_1, m , be integers, $r + r_1 = m + 1$, $r \geq 1, r_1 \geq 1$, and the spline ω_k be such that $\text{supp } \omega_k = [x_{k-r}, x_{k+r_1}]$. Following the methodology developed by Professor S.G. Mikhlin, we find the basis functions by solving the system of approximation relations

$$\sum_{j=k+1-r_1}^{k+r} x_j^s \omega_j(x) = x^s, \quad x \in [x_k, x_{k+1}],$$

$$s = 0, 1, \dots, m. \tag{1}$$

2.1 Polynomial Splines of the Second Order of Approximation

Let $r = 1, r_1 = 1$. The support of the basis splines of the second order of approximation occupies two grid intervals. These splines are convenient to use on a finite interval, both on a uniform grid of nodes and on a non-uniform grid of nodes. The approximation of the function on a finite interval of interpolation does not have a boundary layer. When solving the Fredholm integral equation of the second kind, the minimum number of the grid nodes is two. We set the support of the basis spline as follows: $\text{supp } \omega_j = [x_{j-1}, x_{j+1}]$. On the interval

$[x_j, x_{j+1}]$ we approximate the function $u(x)$ by the following expression:

$$\tilde{u}(x) = u(x_j)\omega_j(x) + u(x_{j+1})\omega_{j+1}(x),$$

$$x \in [x_j, x_{j+1}],$$

where the basis splines $\omega_j(x), \omega_{j+1}(x)$ are as follows:

$$\omega_j(x) = \frac{x - x_{j+1}}{x_j - x_{j+1}}, \quad x \in [x_j, x_{j+1}],$$

$$\omega_{j+1}(x) = \frac{x - x_j}{x_{j+1} - x_j}, \quad x \in [x_j, x_{j+1}].$$

These splines are the interpolation splines of the second order of approximation as well as the first degree. The approximation using these splines is the continuous approximation.

Note that the minimum number of grid intervals is one, and the minimum number of grid nodes is two.

The following statement is valid for the approximation error.

Let $h = x_{j+1} - x_j$. In the case of the splines of the first degree, it is easy to obtain an estimate of the approximation error on the interval $[x_j, x_{j+1}]$, [22], [23]

$$|u(x) - \tilde{u}(x)| \leq \frac{h^2}{8} \max_{[x_j, x_{j+1}]} |u''(x)|,$$

$$x \in [x_j, x_{j+1}].$$

In the case of an uneven grid of nodes, we take the length of the maximum grid interval as the value of h .

Further, we will use the norm of the form:

$$\|u\|_{C[a,b]} = \max_{x \in [a,b]} |u(x)|.$$

Consider the solution of the Fredholm integral equation of the second kind

$$(Au)(x) \equiv u(x) - \int_a^b K(x, s)u(s) ds = f(x),$$

$$x \in [a, b].$$

Here, and further, we assume that the kernel $K(x, s)$ and the right side of the equation $f(x)$ are

continuous. In addition, we assume that the equation is uniquely solvable and the estimate for the norm of the inverse operator in space C is known:

$$\|A^{-1}\| \leq B.$$

Suppose $|K(x, s)| < \rho < 1$, when $0 \leq x \leq 1$, $0 \leq s \leq 1$.

We construct the set of nodes $x_k, k = 0, 1, \dots, n$, on the interval $[a, b]$. We have the relation

$$\int_a^b K(x, s)u(s) ds = \sum_{k=0}^{n-1} \int_{x_k}^{x_{k+1}} K(x, s)u(s) ds.$$

On each interval $[x_k, x_{k+1}]$ we replace $u(s)$ with approximation $\tilde{u}(x)$.

Now we have the integral equation in the form:

$$u(x) - \sum_{k=0}^{n-1} \int_{x_k}^{x_{k+1}} K(x, s)(u_k \omega_k(s) + u_{k+1} \omega_{k+1}(s)) ds = f(x).$$

From here we get the system of equation in the form

$$u(x) - \sum_{k=1}^{n-1} (u_k A_k(x) + u_{k+1} A_{k+1}(x)) = f(x),$$

where

$$A_k(x) = \int_{x_k}^{x_{k+1}} K(x, s) \omega_k(s) ds,$$

$$A_{k+1}(x) = \int_{x_k}^{x_{k+1}} K(x, s) \omega_{k+1}(s) ds.$$

Next, we take x_j instead of x and we have to solve the system of linear algebraic equations:

$$u_j^h - \sum_{k=0}^{n-1} (u_k^h A_k(x_j) + u_{k+1}^h A_{k+1}(x_j)) = f_j^h,$$

$$j = 0, 1, 2, \dots, n - 1.$$

We assume that the integral $\int_{x_k}^{x_{k+1}} K(x, s) \omega_k(s) ds$ can be computed exactly. Otherwise, we can use quadrature formulas.

Now, we suppose that $a = 0, b = 1$, and $h = \text{const}$. Let u_j be one of those components of the solution, whose absolute value is the largest. Therefore, the execution for this component of the solution is $|u_j| = \max_k |u_k|$.

We consider on the interval $[x_k, x_{k+1}]$ the approximation of the function with splines of the second order of approximation

$$\tilde{u}(x) = u_k \omega_k(x) + u_{k+1} \omega_{k+1}(x).$$

Now we have

$$|f(x_j)| = |u(x_j) - \int_0^1 K(x_j, s) u(s) ds|,$$

where

$$\int_0^1 K(x_j, s) u(s) ds \approx \sum_{k=0}^{n-1} \int_{x_k}^{x_{k+1}} K(x_j, s) (u_k \omega_k(s) + u_{k+1} \omega_{k+1}(s)) ds.$$

We assume that $h = x_{k+1} - x_k$. It is easy to calculate the integrals $\int_{x_k}^{x_{k+1}} \omega_k(s) ds = h/2$, $\int_{x_k}^{x_{k+1}} \omega_{k+1}(s) ds = h/2$.

Now using the mean value theorem of integral calculus, we obtain

$$\int_{x_k}^{x_{k+1}} K(x_j, s) \omega_k(s) ds = K(x_j, \eta_k) \int_{x_k}^{x_{k+1}} \omega_k(s) ds = K(x_j, \eta_k) \frac{h}{2}, \eta_k \in [x_k, x_{k+1}].$$

Similarly, we get

$$\int_{x_k}^{x_{k+1}} K(x_j, s) \omega_{k+1}(s) ds = K(x_j, \xi_k) \frac{h}{2}, \xi_k \in [x_k, x_{k+1}].$$

Finally, we obtain the inequality

$$|f(x_j)| \geq (1 - \rho) |u(x_j)|.$$

Thus, we have the estimation

$$|u(x_j)| \leq \frac{1}{1 - \rho} |f(x_j)|.$$

In particular, we can take $f = 0$. This implies that the system is uniquely solvable. The last inequality means the stability of the solution depends on the right side of the equation with the constant $C = \frac{1}{1 - \rho}$.

2.2 Polynomial Splines of the Seventh Order of Approximation

Now we consider the application of splines of the seventh order of approximation to solve the Fredholm integral equations of the second kind. Different modifications of the splines of the seventh order of approximation are used at the beginning, in the middle and at the end of the interpolation interval $[a, b]$. The support of the basis spline occupies seven grid intervals.

First, consider the approximation properties of polynomial splines of the seventh order of approximation.

Let r, r_1 be integers, $r + r_1 = 7, r \geq 1, r_1 \geq 1$, and the spline ω_k be such that $\text{supp } \omega_k =$

$[x_{k-r}, x_{k+r_1}]$. We find the basis functions by solving the system of approximation relations

$$\sum_{j=k+1-r_1}^{k+r} x_j^s \omega_j(x) = x^s, \quad x \in [x_k, x_{k+1}],$$

$$s = 0, 1, \dots, 6.$$

With different values of the parameters r, r_1 , we get basis splines suitable for the approximation at the beginning of the interpolation interval (the right basis splines), in the middle of the interpolation interval (the middle basis splines), or at the end of the interpolation interval (the left basis splines).

With $r_1 = 4$ and $r = 3$ we get the middle splines. On the interval $[x_k, x_{k+1}]$, we construct the approximation with the middle splines at a distance of three grid intervals from the ends of the interval $[a, b]$ in the form:

$$\tilde{u}^M(x) = \sum_{j=k-3}^{k+3} u(x_j) \omega_j^M(x), \quad x \in [x_k, x_{k+1}],$$

where the middle basis splines $\omega_j^M(x)$ have the form:

$$\omega_{k-3}^M(x) = c_{k-3}(x)/d_{k-3},$$

where

$$c_{k-3}(x) = (x - x_{k+3})(x - x_{k+2})(x - x_{k+1}) \times (x - x_k)(x - x_{k-1})(x - x_{k-2}),$$

$$d_{k-3} = (x_{k-3} - x_{k+3})(x_{k-3} - x_{k+2}) \times (x_{k-3} - x_{k+1})(x_{k-3} - x_k) \times (x_{k-3} - x_{k-1})(x_{k-3} - x_{k-2});$$

$$\omega_{k-2}^M(x) = \frac{c_{k-2}(x)}{d_{k-2}},$$

$$c_{k-2}(x) = (x - x_{k+3})(x - x_{k+2})(x - x_{k+1}) \times (x - x_k)(x - x_{k-1})(x - x_{k-3}),$$

$$d_{k-2} = (x_{k-2} - x_{k+3})(x_{k-2} - x_{k+2}) \times (x_{k-2} - x_{k+1})(x_{k-2} - x_k) \times (x_{k-2} - x_{k-1})(x_{k-2} - x_{k-3});$$

$$\omega_{k-1}^M(x) = \frac{c_{k-1}(x)}{d_{k-1}},$$

$$c_{k-1}(x) = (x - x_{k+3})(x - x_{k+2})(x - x_{k+1}) \times (x - x_k)(x - x_{k-2})(x - x_{k-3}),$$

$$d_{k-1} = (x_{k-1} - x_{k+3})(x_{k-1} - x_{k+2}) \times (x_{k-1} - x_{k+1})(x_{k-1} - x_k) \times (x_{k-1} - x_{k-2})(x_{k-1} - x_{k-3});$$

$$\omega_k^M(x) = \frac{c_k(x)}{d_k},$$

$$c_k(x) = (x - x_{k+3})(x - x_{k+2})(x - x_{k+1}) \times (x - x_{k-1})(x - x_{k-2})(x - x_{k-3}),$$

$$d_k = (x_k - x_{k+3})(x_k - x_{k+2})(x_k - x_{k+1})$$

$$\times (x_k - x_{k-1})(x_k - x_{k-2})(x_k - x_{k-3});$$

$$\omega_{k+1}^M(x) = \frac{c_{k+1}(x)}{d_{k+1}},$$

$$c_{k+1}(x) = (x - x_{k+3})(x - x_{k+2})(x - x_k) \times (x - x_{k-1})(x - x_{k-2})(x - x_{k-3}),$$

$$d_{k+1} = (x_{k+1} - x_{k+3})(x_{k+1} - x_{k+2}) \times (x_{k+1} - x_k)(x_{k+1} - x_{k-1}) \times (x_{k+1} - x_{k-2})(x_{k+1} - x_{k-3});$$

$$\omega_{k+2}^M(x) = \frac{c_{k+2}(x)}{d_{k+2}},$$

$$c_{k+2}(x) = (x - x_{k+1})(x - x_k)(x - x_{k-1}) \times (x - x_{k-2})(x - x_{k-3})(x - x_{k+3}),$$

$$d_{k+2} = (x_{k+2} - x_{k+1})(x_{k+2} - x_k) \times (x_{k+2} - x_{k-1})(x_{k+2} - x_{k-2}) \times (x_{k+2} - x_{k-3})(x_{k+2} - x_{k+3}).$$

Approximations with these basis splines can be constructed on the grid intervals $[x_k, x_{k+1}]$, $k = 3, \dots, n - 3$.

Let us consider the approximation with the left basis splines. We get the left basis splines when $r_1 = 6, r = 1$. In this case, formula (1) on the interval $[x_k, x_{k+1}]$ takes the form:

$$\tilde{u}^L(x) = \sum_{j=k-5}^{k+1} u(x_j) \omega_j^L(x), \quad x \in [x_k, x_{k+1}].$$

where the basis splines $\omega_k^L(x)$ have the form

$$\omega_{k-5}^L(x) = c_{k-5}(x)/d_{k-5},$$

$$c_{k-5}(x) = (x - x_{k+1})(x - x_k)(x - x_{k-1}) \times (x - x_{k-2})(x - x_{k-3})(x - x_{k-4}),$$

$$d_{k-5} = (x_{k-5} - x_{k+1})(x_{k-5} - x_k) \times (x_{k-5} - x_{k-1})(x_{k-5} - x_{k-2}) \times (x_{k-5} - x_{k-3})(x_{k-5} - x_{k-4});$$

$$\omega_{k-4}^L(x) = \frac{c_{k-4}(x)}{d_{k-4}},$$

$$c_{k-4}(x) = (x - x_{k+1})(x - x_k)(x - x_{k-1}) \times (x - x_{k-2})(x - x_{k-3})(x - x_{k-5}),$$

$$d_{k-4} = (x_{k-4} - x_{k+1})(x_{k-4} - x_k) \times (x_{k-4} - x_{k-1})(x_{k-4} - x_{k-2}) \times (x_{k-4} - x_{k-3})(x_{k-4} - x_{k-5});$$

$$\omega_{k-3}^L(x) = \frac{c_{k-3}(x)}{d_{k-3}},$$

$$c_{k-3}(x) = (x - x_{k+1})(x - x_k)(x - x_{k-1}) \times (x - x_{k-2})(x - x_{k-4})(x - x_{k-5}),$$

$$d_{k-3} = (x_{k-3} - x_{k+1})(x_{k-3} - x_k)$$

$$\begin{aligned} &\times (x_{k-3} - x_{k-1})(x_{k-3} - x_{k-2}) \\ &\times (x_{k-3} - x_{k-4})(x_{k-3} - x_{k-5}); \end{aligned}$$

$$\omega_{k-2}^L(x) = \frac{c_{k-2}(x)}{d_{k-2}},$$

$$\begin{aligned} c_{k-2}(x) &= (x - x_{k+1})(x - x_k)(x - x_{k-1}) \\ &\times (x - x_{k-3})(x - x_{k-4})(x - x_{k-5}), \\ d_{k-2} &= (x_{k-2} - x_{k+1})(x_{k-2} - x_k) \\ &\times (x_{k-2} - x_{k-1})(x_{k-2} - x_{k-3}) \\ &\times (x_{k-2} - x_{k-4})(x_{k-2} - x_{k-5}); \end{aligned}$$

$$\omega_{k-1}^L(x) = \frac{c_{k-1}(x)}{d_{k-1}},$$

$$\begin{aligned} c_{k-1}(x) &= (x - x_{k+1})(x - x_k)(x - x_{k-2}) \\ &\times (x - x_{k-3})(x - x_{k-4})(x - x_{k-5}), \\ d_{k-1} &= (x_{k-1} - x_{k+1})(x_{k-1} - x_k) \\ &\times (x_{k-1} - x_{k-2})(x_{k-1} - x_{k-3}) \\ &\times (x_{k-1} - x_{k-4})(x_{k-1} - x_{k-5}); \end{aligned}$$

$$\omega_k^L(x) = \frac{c_k(x)}{d_k},$$

$$\begin{aligned} c_k(x) &= (x - x_{k+1})(x - x_{k-1})(x - x_{k-2}) \\ &(x - x_{k-3})(x - x_{k-4})(x - x_{k-5}), \\ d_k &= (x_k - x_{k+1})(x_k - x_{k-1})(x_k - x_{k-2}) \\ &(x_k - x_{k-3})(x_k - x_{k-4})(x_k - x_{k-5}); \end{aligned}$$

$$\omega_{k+1}^L(x) = \frac{c_{k+1}(x)}{d_{k+1}},$$

$$\begin{aligned} c_{k+1}(x) &= (x - x_k)(x - x_{k-1})(x - x_{k-2}) \\ &\times (x - x_{k-3})(x - x_{k-4})(x - x_{k-5}), \\ d_{k+1} &= (x_{k+1} - x_k)(x_{k+1} - x_{k-1}) \\ &\times (x_{k+1} - x_{k-2})(x_{k+1} - x_{k-3}) \\ &\times (x_{k+1} - x_{k-4})(x_{k+1} - x_{k-5}). \end{aligned}$$

Approximations with these basis splines can be applied on the next grid interval $[x_k, x_{k+1}]$, $k = 5, \dots, n - 1$.

Let us consider the approximation with the left-right basis splines. We get the left basis splines when $r_1 = 5$, $r = 2$. In this case, formula (1) on the interval $[x_k, x_{k+1}]$ takes the form

$$\tilde{u}^{LR}(x) = \sum_{j=k-4}^{k+2} u(x_j)\omega_j^{LR}(x), x \in [x_k, x_{k+1}],$$

where the basis splines $\omega_k^{LR}(x)$ have the form

$$\omega_{k-4}^{LR}(x) = c_{k-4}(x)/d_{k-4},$$

$$\begin{aligned} c_{k-4}(x) &= (x - x_{k+2})(x - x_{k+1})(x - x_k) \\ &\times (x - x_{k-1})(x - x_{k-2})(x - x_{k-3}), \\ d_{k-4} &= (x_{k-4} - x_{k+2})(x_{k-4} - x_{k+1}) \end{aligned}$$

$$\begin{aligned} &\times (x_{k-4} - x_k)(x_{k-4} - x_{k-1}) \\ &\times (x_{k-4} - x_{k-2})(x_{k-4} - x_{k-3}); \end{aligned}$$

$$\omega_{k-3}^{LR}(x) = \frac{c_{k-3}(x)}{d_{k-3}},$$

$$\begin{aligned} c_{k-3}(x) &= (x - x_{k+2})(x - x_{k+1})(x - x_k) \\ &\times (x - x_{k-1})(x - x_{k-2})(x - x_{k-4}), \\ d_{k-3} &= (x_{k-3} - x_{k+2})(x_{k-3} - x_{k+1}) \\ &\times (x_{k-3} - x_k)(x_{k-3} - x_{k-1}) \\ &\times (x_{k-3} - x_{k-2})(x_{k-3} - x_{k-4}); \end{aligned}$$

$$\omega_{k-2}^{LR}(x) = \frac{c_{k-2}(x)}{d_{k-2}},$$

$$\begin{aligned} c_{k-2}(x) &= (x - x_{k+2})(x - x_{k+1})(x - x_k) \\ &\times (x - x_{k-1})(x - x_{k-3})(x - x_{k-4}), \\ d_{k-2} &= (x_{k-2} - x_{k+2})(x_{k-2} - x_{k+1}) \\ &\times (x_{k-2} - x_k)(x_{k-2} - x_{k-1}) \\ &\times (x_{k-2} - x_{k-3})(x_{k-2} - x_{k-4}); \end{aligned}$$

$$\omega_{k-1}^{LR}(x) = \frac{c_{k-1}(x)}{d_{k-1}},$$

$$\begin{aligned} c_{k-1}(x) &= (x - x_{k+2})(x - x_{k+1})(x - x_k) \\ &\times (x - x_{k-2})(x - x_{k-3})(x - x_{k-4}), \\ d_{k-1} &= (x_{k-1} - x_{k+2})(x_{k-1} - x_{k+1}) \\ &\times (x_{k-1} - x_k)(x_{k-1} - x_{k-2}) \\ &\times (x_{k-1} - x_{k-3})(x_{k-1} - x_{k-4}); \end{aligned}$$

$$\omega_k^{LR}(x) = \frac{c_k(x)}{d_k},$$

$$\begin{aligned} c_k(x) &= (x - x_{k+2})(x - x_{k+1})(x - x_{k-1}) \\ &\times (x - x_{k-2})(x - x_{k-3})(x - x_{k-4}), \\ d_k &= (x_k - x_{k+2})(x_k - x_{k+1}) \\ &\times (x_k - x_{k-1})(x_k - x_{k-2}) \\ &\times (x_k - x_{k-3})(x_k - x_{k-4}); \end{aligned}$$

$$\omega_{k+1}^{LR}(x) = \frac{c_{k+1}(x)}{d_{k+1}},$$

$$\begin{aligned} c_{k+1}(x) &= (x - x_{k+2})(x - x_k)(x - x_{k-1}) \\ &\times (x - x_{k-2})(x - x_{k-3})(x - x_{k-4}), \\ d_{k+1} &= (x_{k+1} - x_{k+2})(x_{k+1} - x_k) \\ &\times (x_{k+1} - x_{k-1})(x_{k+1} - x_{k-2}) \\ &\times (x_{k+1} - x_{k-3})(x_{k+1} - x_{k-4}); \end{aligned}$$

$$\omega_{k+2}^{LR}(x) = \frac{c_{k+2}(x)}{d_{k+2}},$$

$$\begin{aligned} c_{k+2}(x) &= (x - x_{k+1})(x - x_k)(x - x_{k-1}) \\ &\times (x - x_{k-2})(x - x_{k-3})(x - x_{k-4}), \\ d_{k+2} &= (x_{k+2} - x_{k-1})(x_{k+2} - x_k) \\ &\times (x_{k+2} - x_{k-1})(x_{k+2} - x_{k-2}) \\ &\times (x_{k+2} - x_{k-3})(x_{k+2} - x_{k-4}). \end{aligned}$$

Approximations with these basis splines can be applied on the next grid intervals $[x_k, x_{k+1}]$, $k = 4, \dots, n - 2$.

Consider the approximation with the right basis splines. Let $r_1 = 1$, $r = 6$. In this case, formula (1) takes the next form on the interval $[x_k, x_{k+1}]$:

$$\tilde{u}^R(x) = \sum_{j=k}^{k+6} u(x_j) \omega_j^R(x), \quad x \in [x_k, x_{k+1}],$$

where the right basis splines $\omega_j^R(x)$ have the form:

$$\omega_k^R(x) = \frac{c_k(x)}{d_k},$$

$$c_k(x) = (x - x_{k+6})(x - x_{k+5})(x - x_{k+4}) \times (x - x_{k+3})(x - x_{k+2})(x - x_{k+1}),$$

$$d_k = (x_k - x_{k+6})(x_k - x_{k+5})(x_k - x_{k+4}) \times (x_k - x_{k+3})(x_k - x_{k+2})(x_k - x_{k+1});$$

$$\omega_{k+1}^R(x) = \frac{c_{k+1}(x)}{d_{k+1}},$$

$$c_{k+1}(x) = (x - x_{k+6})(x - x_{k+5})(x - x_{k+4}) \times (x - x_{k+3})(x - x_{k+2})(x - x_k),$$

$$d_{k+1} = (x_{k+1} - x_{k+6})(x_{k+1} - x_{k+5}) \times (x_{k+1} - x_{k+4})(x_{k+1} - x_{k+3}) \times (x_{k+1} - x_{k+2})(x_{k+1} - x_k);$$

$$\omega_{k+2}^R(x) = \frac{c_{k+2}(x)}{d_{k+2}},$$

$$c_{k+2}(x) = (x - x_{k+6})(x - x_{k+5})(x - x_{k+4}) \times (x - x_{k+3})(x - x_{k+1})(x - x_k),$$

$$d_{k+2} = (x_{k+2} - x_{k+6})(x_{k+2} - x_{k+5}) \times (x_{k+2} - x_{k+4})(x_{k+2} - x_{k+3}) \times (x_{k+2} - x_{k+1})(x_{k+2} - x_k);$$

$$\omega_{k+3}^R(x) = \frac{c_{k+3}(x)}{d_{k+3}},$$

$$c_{k+3}(x) = (x - x_{k+6})(x - x_{k+5})(x - x_{k+4}) \times (x - x_{k+2})(x - x_{k+1})(x - x_k),$$

$$d_{k+3} = (x_{k+3} - x_{k+6})(x_{k+3} - x_{k+5}) \times (x_{k+3} - x_{k+4})(x_{k+3} - x_{k+2}) \times (x_{k+3} - x_{k+1})(x_{k+3} - x_k);$$

$$\omega_{k+4}^R(x) = \frac{c_{k+4}(x)}{d_{k+4}},$$

$$c_{k+4}(x) = (x - x_{k+6})(x - x_{k+5})(x - x_{k+3}) \times (x - x_{k+2})(x - x_{k+1})(x - x_k),$$

$$d_{k+4} = (x_{k+4} - x_{k+6})(x_{k+4} - x_{k+5}) \times (x_{k+4} - x_{k+3})(x_{k+4} - x_{k+2}) \times (x_{k+4} - x_{k+1})(x_{k+4} - x_k);$$

$$\omega_{k+5}^R(x) = \frac{c_{k+5}(x)}{d_{k+5}},$$

$$c_{k+5}(x) = (x - x_{k+6})(x - x_{k+4})(x - x_{k+3}) \times (x - x_{k+2})(x - x_{k+1})(x - x_k),$$

$$d_{k+5} = (x_{k+5} - x_{k+6})(x_{k+5} - x_{k+4}) \times (x_{k+5} - x_{k+3})(x_{k+5} - x_{k+2}) \times (x_{k+5} - x_{k+1})(x_{k+5} - x_k);$$

$$\omega_{k+6}^R(x) = \frac{c_{k+6}(x)}{d_{k+6}},$$

$$c_{k+6}(x) = (x - x_{k+5})(x - x_{k+4})(x - x_{k+3}) \times (x - x_{k+2})(x - x_{k+1})(x - x_k),$$

$$d_{k+6} = (x_{k+6} - x_{k+5})(x_{k+6} - x_{k+4}) \times (x_{k+6} - x_{k+3})(x_{k+6} - x_{k+2}) \times (x_{k+6} - x_{k+1})(x_{k+6} - x_k).$$

Approximations with these basis splines can be applied on the next grid intervals $[x_k, x_{k+1}]$, $k = 0, \dots, n - 6$.

Consider the approximation with the right-left basis splines. Let $r_1 = 2$, $r = 5$, in this case, on the interval $[x_k, x_{k+1}]$ formula (1) takes the form:

$$\tilde{u}^{RL}(x) = \sum_{j=k-1}^{k+5} u(x_j) \omega_j^{RL}(x), \quad x \in [x_k, x_{k+1}],$$

where the right-left basis splines $\omega_j^R(x)$ have the form:

$$\omega_{k+5}^{RL}(x) = \frac{c_{k+5}(x)}{d_{k+5}},$$

$$c_{k+5}(x) = (x - x_{k+4})(x - x_{k+3})(x - x_{k+2}) \times (x - x_{k+1})(x - x_k)(x - x_{k-1}),$$

$$d_{k+5} = (x_{k+5} - x_{k+4})(x_{k+5} - x_{k+3}) \times (x_{k+5} - x_{k+2})(x_{k+5} - x_{k+1}) \times (x_{k+5} - x_k)(x_{k+5} - x_{k-1});$$

$$\omega_{k+4}^{RL}(x) = \frac{c_{k+4}(x)}{d_{k+4}},$$

$$c_{k+4}(x) = (x - x_{k+5})(x - x_{k+3})(x - x_{k+2}) \times (x - x_{k+1})(x - x_k)(x - x_{k-1}),$$

$$d_{k+4} = (x_{k+4} - x_{k+5})(x_{k+4} - x_{k+3}) \times (x_{k+4} - x_{k+2})(x_{k+4} - x_{k+1}) \times (x_{k+4} - x_k)(x_{k+4} - x_{k-1});$$

$$\omega_{k+3}^{RL}(x) = \frac{c_{k+3}(x)}{d_{k+3}},$$

$$c_{k+3}(x) = (x - x_{k+5})(x - x_{k+4})(x - x_{k+2}) \times (x - x_{k+1})(x - x_k)(x - x_{k-1}),$$

$$d_{k+3} = (x_{k+3} - x_{k+5})(x_{k+3} - x_{k+4}) \times (x_{k+3} - x_{k+2})(x_{k+3} - x_{k+1}) \times (x_{k+3} - x_k)(x_{k+3} - x_{k-1});$$

$$\omega_{k+2}^{RL}(x) = \frac{c_{k+2}(x)}{d_{k+2}},$$

$$\begin{aligned} c_{k+2}(x) &= (x - x_{k+5})(x - x_{k+4})(x - x_{k+3}) \\ &\times (x - x_{k+1})(x - x_k)(x - x_{k-1}), \\ d_{k+2} &= (x_{k+2} - x_{k+5})(x_{k+2} - x_{k+4}) \\ &\times (x_{k+2} - x_{k+3})(x_{k+2} - x_{k+1}) \\ &\times (x_{k+2} - x_k)(x_{k+2} - x_{k-1}); \end{aligned}$$

$$\omega_{k+1}^{RL}(x) = \frac{c_{k+1}(x)}{d_{k+1}},$$

$$\begin{aligned} c_{k+1}(x) &= (x - x_{k+5})(x - x_{k+4})(x - x_{k+3}) \\ &\times (x - x_{k+2})(x - x_k)(x - x_{k-1}), \\ d_{k+1} &= (x_{k+1} - x_{k+5})(x_{k+1} - x_{k+4}) \\ &\times (x_{k+1} - x_{k+3})(x_{k+1} - x_{k+2}) \\ &\times (x_{k+1} - x_k)(x_{k+1} - x_{k-1}); \end{aligned}$$

$$\omega_k^{RL}(x) = \frac{c_k(x)}{d_k},$$

$$\begin{aligned} c_k(x) &= (x - x_{k+5})(x - x_{k+4})(x - x_{k+3}) \\ &\times (x - x_{k+2})(x - x_{k+1})(x - x_{k-1}), \\ d_k &= (x_k - x_{k+5})(x_k - x_{k+4}) \\ &\times (x_k - x_{k+3})(x_k - x_{k+2}) \\ &\times (x_k - x_{k+1})(x_k - x_{k-1}); \end{aligned}$$

$$\omega_{k-1}^{RL}(x) = \frac{c_{k-1}(x)}{d_{k-1}},$$

$$\begin{aligned} c_{k-1}(x) &= (x - x_{k+5})(x - x_{k+4})(x - x_{k+3}) \\ &\times (x - x_{k+2})(x - x_{k+1})(x - x_k), \\ d_{k-1} &= (x_{k-1} - x_{k+5})(x_{k-1} - x_{k+4}) \\ &\times (x_{k-1} - x_{k+3})(x_{k-1} - x_{k+2}) \\ &\times (x_{k-1} - x_{k+1})(x_{k-1} - x_k). \end{aligned}$$

Approximations with these basis splines can be applied on the next grid intervals $[x_k, x_{k+1}]$, $k = 1, \dots, n - 5$.

Consider the approximation with the right-left-left basis splines. Let $r_1 = 3$, $r = 4$. In this case, on the interval $[x_k, x_{k+1}]$ formula (1) takes the form:

$$\begin{aligned} \tilde{u}^{RLL}(x) &= \sum_{j=k-2}^{k+4} u(x_j)\omega_j^{RLL}(x), \\ x &\in [x_k, x_{k+1}], \end{aligned}$$

where the right-left-left basis splines $\omega_j^{RLL}(x)$ have the form:

$$\omega_{k+4}^{RLL}(x) = \frac{c_{k+4}(x)}{d_{k+4}},$$

$$\begin{aligned} c_{k+4}(x) &= (x - x_{k+3})(x - x_{k+2})(x - x_{k+1}) \\ &\times (x - x_k)(x - x_{k-1})(x - x_{k-2}), \\ d_{k+4} &= (x_{k+4} - x_{k+3})(x_{k+4} - x_{k+2}) \end{aligned}$$

$$\begin{aligned} &\times (x_{k+4} - x_{k+1})(x_{k+4} - x_k) \\ &\times (x_{k+4} - x_{k-1})(x_{k+4} - x_{k-2}); \end{aligned}$$

$$\omega_{k+3}^{RLL}(x) = \frac{c_{k+3}(x)}{d_{k+3}},$$

$$\begin{aligned} c_{k+3}(x) &= (x - x_{k+4})(x - x_{k+2})(x - x_{k+1}) \\ &\times (x - x_k)(x - x_{k-1})(x - x_{k-2}), \\ d_{k+3} &= (x_{k+3} - x_{k+4})(x_{k+3} - x_{k+2}) \\ &\times (x_{k+3} - x_{k+1})(x_{k+3} - x_k) \\ &\times (x_{k+3} - x_{k-1})(x_{k+3} - x_{k-2}); \end{aligned}$$

$$\omega_{k+2}^{RLL}(x) = \frac{c_{k+2}(x)}{d_{k+2}},$$

$$\begin{aligned} c_{k+2}(x) &= (x - x_{k+4})(x - x_{k+3})(x - x_{k+1}) \\ &\times (x - x_k)(x - x_{k-1})(x - x_{k-2}), \\ d_{k+2} &= (x_{k+2} - x_{k+4})(x_{k+2} - x_{k+3}) \\ &\times (x_{k+2} - x_{k+1})(x_{k+2} - x_k) \\ &\times (x_{k+2} - x_{k-1})(x_{k+2} - x_{k-2}); \end{aligned}$$

$$\omega_{k+1}^{RLL}(x) = \frac{c_{k+1}(x)}{d_{k+1}},$$

$$\begin{aligned} c_{k+1}(x) &= (x - x_{k+4})(x - x_{k+3})(x - x_{k+2}) \\ &\times (x - x_k)(x - x_{k-1})(x - x_{k-2}), \\ d_{k+1} &= (x_{k+1} - x_{k+4})(x_{k+1} - x_{k+3}) \\ &\times (x_{k+1} - x_{k+2})(x_{k+1} - x_k) \\ &\times (x_{k+1} - x_{k-1})(x_{k+1} - x_{k-2}); \end{aligned}$$

$$\omega_k^{RLL}(x) = \frac{c_k(x)}{d_k},$$

$$\begin{aligned} c_k(x) &= (x - x_{k+4})(x - x_{k+3})(x - x_{k+2}) \\ &\times (x - x_{k+1})(x - x_{k-1})(x - x_{k-2}), \\ d_k &= (x_k - x_{k+4})(x_k - x_{k+3}) \\ &\times (x_k - x_{k+2})(x_k - x_{k+1}) \\ &\times (x_k - x_{k-1})(x_k - x_{k-2}); \end{aligned}$$

$$\omega_{k-1}^{RLL}(x) = \frac{c_{k-1}(x)}{d_{k-1}},$$

$$\begin{aligned} c_{k-1}(x) &= (x - x_{k+4})(x - x_{k+3})(x - x_{k+2}) \\ &\times (x - x_{k-1})(x - x_k)(x - x_{k-2}), \\ d_{k-1} &= (x_{k-1} - x_{k+4})(x_{k-1} - x_{k+3}) \\ &\times (x_{k-1} - x_{k+2})(x_{k-1} - x_{k+1}) \\ &\times (x_{k-1} - x_k)(x_{k-1} - x_{k-2}); \end{aligned}$$

$$\omega_{k-2}^{RLL}(x) = \frac{c_{k-2}(x)}{d_{k-2}},$$

$$\begin{aligned} c_{k-2}(x) &= (x - x_{k+4})(x - x_{k+3})(x - x_{k+2}) \\ &\times (x - x_{k+1})(x - x_k)(x - x_{k-1}), \\ d_{k-2} &= (x_{k-2} - x_{k+4})(x_{k-2} - x_{k+3}) \\ &\times (x_{k-2} - x_{k+2})(x_{k-2} - x_{k+1}) \\ &\times (x_{k-2} - x_k)(x_{k-2} - x_{k-1}). \end{aligned}$$

Approximations with these basis splines can be applied on the next grid intervals $[x_k, x_{k+1}]$, $k = 2, \dots, n - 4$.

When approximating a function with the splines of the 7th order of approximation, the next Theorem is valid.

Theorem. If $\text{supp } \omega_k = [x_{k-1}, x_{k+6}]$, then the following inequality is valid:

$$|u(x) - \tilde{u}^L(x)|_{x \in [x_k, x_{k+1}]} \leq h^7 \frac{95.842}{7!} \|u^{(7)}\|_{C[x_{k-5}, x_{k+1}]}.$$

If $\text{supp } \omega_k = [x_{k-2}, x_{k+5}]$, then the following inequality is valid:

$$|u(x) - \tilde{u}^{LR}(x)|_{x \in [x_k, x_{k+1}]} \leq h^7 \frac{23.149}{7!} \|u^{(7)}\|_{C[x_{k-4}, x_{k+2}]}.$$

If $\text{supp } \omega_k = [x_{k-3}, x_{k+4}]$, then the following approximation estimate is valid:

$$|u(x) - \tilde{u}^M(x)|_{x \in [x_k, x_{k+1}]} \leq h^7 \frac{12.359}{7!} \|u^{(7)}\|_{C[x_{k-3}, x_{k+3}]}.$$

If $\text{supp } \omega_k = [x_{k-4}, x_{k+3}]$, then the following approximation estimate is valid:

$$|u(x) - \tilde{u}^{RLL}(x)|_{x \in [x_k, x_{k+1}]} \leq h^7 \frac{12.359}{7!} \|u^{(7)}\|_{C[x_{k-2}, x_{k+4}]}.$$

If $\text{supp } \omega_k = [x_{k-5}, x_{k+2}]$, then the following approximation estimate is valid:

$$|u(x) - \tilde{u}^{RL}(x)|_{x \in [x_k, x_{k+1}]} \leq h^7 \frac{23.149}{7!} \|u^{(7)}\|_{C[x_{k-1}, x_{k+5}]}.$$

If $\text{supp } \omega_k = [x_{k-6}, x_{k+1}]$, then the following approximation estimate is valid:

$$|u(x) - \tilde{u}^R(x)|_{x \in [x_k, x_{k+1}]} \leq h^7 \frac{95.842}{7!} \|u^{(7)}\|_{C[x_k, x_{k+6}]}.$$

Proof. In the case of approximating the function u on the interval $[x_k, x_{k+1}]$ near the left end of the interval $[a, b]$, we use the right basis splines

$$\tilde{u}(x) = \sum_{j=k}^{k+6} u(x_j) \omega_j^R(x) dx, x \in [x_k, x_{k+1}].$$

First, we estimate the approximation error on the interval $[x_k, x_{k+1}]$ when the right basis splines are used. Using the formula of the remainder term of the interpolation polynomial that solves the Lagrange interpolation problem, we obtain the relation

$$u(x) - \tilde{u}(x) = \frac{1}{7!} (x - x_k) \dots (x - x_{k+6}) u^{(7)}(\xi), \quad \xi \in [x_{k-5}, x_{k+1}].$$

There is a product $(x - x_k) \dots (x - x_{k+6})$ in the error estimate. Let the ordered grid of nodes $\{x_k\}$ be uniform with step h . Let us estimate the product of factors $(x - x_k) \dots (x - x_{k+6})$.

Thus, estimating the maximum of the expression $\frac{1}{7!} (x - x_k) \dots (x - x_{k+6}) u^{(7)}(\xi)$, where $\xi \in [x_k, x_{k+6}]$, we obtain

$$\|u(x) - \tilde{u}(x)\|_{C[x_k, x_{k+1}]} \leq K h^7 \|u^{(7)}\|_{C[x_k, x_{k+6}]}.$$

Similarly, we obtain an approximation estimate on the grid interval $[x_k, x_{k+1}]$ with the left and middle splines.

This completes the proof of the theorem.

Remark 1. The approximation on the interval.

If the interval $[a, b]$ is divided into 6 grid intervals, then it is possible to construct an approximation on the entire interval using the previously presented approximations on the intervals $[x_k, x_{k+1}]$, as follows:

for $k = 0$ we apply the approximation $\tilde{u}^R(x)$;
 for $k = 1$ we apply the approximation $\tilde{u}^{RL}(x)$;
 for $k = 2$ we apply the approximation $\tilde{u}^{RLL}(x)$;
 for $k = 3$ we apply the approximation $\tilde{u}^M(x)$;
 for $k = 4$ we apply the approximation $\tilde{u}^{LR}(x)$;
 for $k = 5$ we apply the approximation $\tilde{u}^L(x)$.

Remark 2. The stability.

Let $a = 0$, $b = 1$, and $x_{k+1} - x_k = h = \text{const}$. Suppose $|K(x, s)| < \rho < 1$, when $0 \leq x \leq 1$, $0 \leq s \leq 1$. The constant in the stability inequality in the case of splines of the seventh order of approximation is calculated in the same way as it was done for the splines of the second order of approximation. To calculate the constant in the stability inequality, we need the following values:

$$\sum_{j=k-2}^{k+4} \int_{x_k}^{x_{k+1}} |\omega_j^{RLL}(x)| dx \approx 1.321 h,$$

$$\begin{aligned} \sum_{j=k-1}^{k+5} \int_{x_k}^{x_{k+1}} |\omega_j^{RL}(x)| dx &\approx 1.635 h, \\ \sum_{j=k}^{k+6} \int_{x_k}^{x_{k+1}} |\omega_j^R(x)| dx &\approx 3.233 h, \\ \sum_{j=k-3}^{k+3} \int_{x_k}^{x_{k+1}} |\omega_j^M(x)| dx &\approx 1.321 h, \\ \sum_{j=k-4}^{k+2} \int_{x_k}^{x_{k+1}} |\omega_j^{LR}(x)| dx &\approx 1.635h, \\ \sum_{j=k-5}^{k+1} \int_{x_k}^{x_{k+1}} |\omega_j^L(x)| dx &\approx 3.233 h. \end{aligned}$$

Taking into account the inequalities given above, we obtain the constants

$$\begin{aligned} s_1 &= (2 \cdot 3.233h + (n - 2) \cdot 1.321h)/n, \\ s_2 &= (3.233h + 1.635h + 1.321h) \cdot 2 \\ &+ (n - 6) \cdot 1.321/n. \end{aligned}$$

Thus, we have the estimations

$$|u(x_j)| \leq \frac{1}{1 - \rho s_i} |f(x_j)|, \quad j = 0, \dots, n, i = 1, 2.$$

Suppose that $|\rho s_i| < 1$. In this case, taking into account the approximation theorem and the inequalities given above, we can see that the approximate solution obtained with the splines of the seventh order of approximation tends to the solution to the Fredholm equation.

3 Problem Solution

In this section, we discuss the solution of the integral equation of the second kind

$$Au \equiv u(x) - \int_a^b K(x, s)u(s)ds = f(x), \quad x \in [a, b],$$

with the local splines of the seventh order of approximation.

Let us choose an integer $n \geq 7$. We build a grid of nodes $\{s_j\}$.

The function $g(s) = K(x, s)u(s), s \in [s_k, s_{k+1}]$, can be approximated with the expression: $g(s) \approx \tilde{g}(s) = K(x, s)\tilde{u}(s)$. Let us denote $c_j = u(s_j)$.

We represent the integral in the form

$$\int_a^b g(s)ds = \sum_{k=0}^{n-1} \int_{s_k}^{s_{k+1}} g(s)ds.$$

Using the results from the second section, we can reduce the integral equation to the solution of a system of linear algebraic equations. To do this, we put $x = x_m, m = 0, \dots, n - 1$, in the equation

$$\begin{aligned} u(x) - \sum_{k=0}^2 \sum_{j=k}^{k+6} c_j \int_{s_k}^{s_{k+1}} K(x, s)\omega_j^R(s)ds \\ - \sum_{k=3}^{n-3} \sum_{j=k-3}^{k+3} c_j \int_{s_k}^{s_{k+1}} K(x, s)\omega_j^M(s)ds \\ - \sum_{k=n-2}^{n-1} \sum_{j=n-3}^{n-1} c_j \int_{s_k}^{s_{k+1}} K(x, s)\omega_j^L(s)ds = f(x). \end{aligned}$$

And now we have to solve the system of linear algebraic equations

$$\begin{aligned} u(x_m) - \sum_{k=0}^2 \sum_{j=k}^{k+6} c_j \int_{s_k}^{s_{k+1}} K(x_m, s)\omega_j^R(s)ds + \\ \sum_{k=3}^{n-2} \sum_{j=k-3}^{k+3} c_j \int_{s_k}^{s_{k+1}} K(x_m, s)\omega_j^M(s)ds + \\ \sum_{k=n-2}^{n-1} \sum_{j=n-3}^{n-1} c_j \int_{s_k}^{s_{k+1}} K(x_m, s)\omega_j^L(s)ds = f(x_m), \\ m = 0, \dots, n. \end{aligned}$$

We can also apply a more detailed approximation (as shown in sections 2.2.1-2.2.6) on the grid intervals adjacent to the boundaries of the interval $[a, b]$. We assume that the integral $\int_{s_k}^{s_{k+1}} v(s)ds = \int_{s_k}^{s_{k+1}} K(x, s)\omega_j(s)ds$ can be computed exactly. Otherwise, we can use quadrature formulas. In this case, it is necessary to take into account the error of the applied quadrature formula. We can use, for example, the Simpson's compound formula:

$$\begin{aligned} \int_{s_k}^{s_{k+1}} v(x)dx = \\ = \frac{(s_{k+1} - s_k)}{3N} (v_0 + 4(v_1 + v_3 + \dots + v_{N-1}) \\ + 2(v_2 + v_4 + \dots + v_{N-2}) + v_N) + R_N(v). \end{aligned}$$

Here N is even, $N \geq 2$.

The formula for the remainder of the Simpson's compound quadrature rule is well known. In the case of the interval $[s_k, s_{k+1}]$, it has the form

$$R_n(v) = \frac{-(s_{k+1} - s_k)^5}{180N^4} v^{(IV)}(\xi),$$

$$\xi \in [s_k, s_{k+1}].$$

Proceeding from this formula and taking into account the approximation theorem for the splines of the seventh order, it is easy to get a priori estimation of the number of nodes of the Simpson's quadrature formula.

Let us calculate the integral from the Runge function $\int_0^{0.4} v(x)dx$, where $v(x) = \frac{1}{1+25x^2}$.

We can obtain

$$\max_{x \in [0, 0.4]} |v^{(IV)}(x)| \leq 15000,$$

$$\max_{x \in [0, 0.4]} |v^{(VII)}(x)| < 0.344 \cdot 10^9.$$

We can easily calculate $N = 2$, and $|R_2(v)| \leq 0.46 \cdot 10^{-6}$. Note that this number is less than the theoretical error of integration (which follows from the Theorem). We can easily obtain that the theoretical error of the integration is about $0.43 \cdot 10^{-4}$.

After the approximate solution at the grid nodes is obtained, we can use the expressions

$$u(x) = \sum_{k=0}^2 \sum_{j=k}^{k+6} c_j \int_{s_k}^{s_{k+1}} K(x, s) \omega_j^R(s) ds$$

$$+ \sum_{k=3}^{n-3} \sum_{j=k-3}^{k+3} c_j \int_{s_k}^{s_{k+1}} K(x, s) \omega_j^M(s) ds$$

$$+ \sum_{k=n-2}^{n-1} \sum_{j=n-3}^{n-1} c_j \int_{s_k}^{s_{k+1}} K(x, s) \omega_j^L(s) ds + f(x).$$

to solve the problems that arise further, for example, the construction of a plot of the solution.

Let us consider two examples of the application of splines of the seventh order of approximation and splines of the second order of approximation to solve the Fredholm integral equation of the second kind.

Example 1. Consider the equation

$$u(x) - 0.1 \int_0^1 \sin\left(\frac{xs}{5}\right) u(s) ds = x \exp(x),$$

$$x \in [0, 1].$$

First, let us construct an ordered grid of nodes x_j with step $h = 1/n$ ($n = 10$) on the interval $[a, b]$. Using splines of the seventh order of approximation, we calculate the approximate values of the solution at these nodes. Next, we construct a sequence of refining grids on the interval $[a, b]$ as follows: we divide each grid interval in half. The division points are added to the nodes of the previous grid.

Thus, we get a new grid of nodes. We compare the values of the approximate solution at some grid nodes.

Table 1 presents the approximate values of the solution at some grid nodes when using splines of the seventh order of approximation when $n = 10, 20$. Table 2 presents the approximate values of the solution at the same grid nodes when using splines of the second order of approximation when $n = 10, 20, 50$.

Table 1. The Approximate Values of the Solution when Splines of the Seventh Order of Approximation were used

x_j	Splines of the Seventh Order of Approximation	
	$n = 10$	$n = 20$
0	0	0
0.2	0.24717242	0.24717241
0.4	0.60251062	0.60251061
0.6	1.10193492	1.10193491
0.8	1.79197032	1.79197030
1.0	2.73268142	2.73268140

Table 2. The Approximate Values of the Solution when Splines of the Second Order of Approximation were used

x_j	Splines of the Second Order of Approximation		
	$n = 10$	$n = 20$	$n = 50$
0	0	0	0
0.2	0.24717235	0.24717240	0.24717241
0.4	0.60251051	0.60251059	0.60251061
0.6	1.10193482	1.10193489	1.10193490
0.8	1.79197029	1.79197031	1.79197030
1.0	2.73268156	2.73268145	2.73268140

Having solved the system of linear algebraic equations, we obtain the framework of an approximate solution. The obtained values of the approximate solution at the grid nodes are marked with circles. Further, using the integral equation and

the found approximate values of the function, we connect the found values with a line. The Maple package was used in the calculations. The plot of the approximate solution with the splines of the second order of approximation when $n = 10$ is given in Figure 1. The nodes are marked along the abscissa axis.




Fig. 1: The plot of the approximate solution when $n = 10$.

Example 2. Consider the equation

$$u(x) - 0.7 \int_0^1 \sin(x + s/5) u(s) ds = \sin(x).$$

As is known, in order to obtain an approximate solution of the integral equation, it is necessary to apply different methods and calculate with a different number of grid nodes. Let us start with calculations using splines of the second order of approximation. Table 3 presents the approximate values of the solution at grid nodes when using splines of the second order of approximation when $n = 10, 20$. Table 4 presents the approximate values of the solution at the same grid nodes when using splines of the second order of approximation when $n = 40, 80$. Table 5 presents the approximate values of the solution at the grid nodes when splines of the second order of approximation when $n = 640$ and splines of the seventh order of approximation when $n = 640$ were used. Table 6 shows the approximate values of the solution when splines of the seventh order of approximation were used ($n = 20, n = 40$).

Table 3. The Approximate Values of the Solution when Splines of the Second Order of Approximation were used

x_j	Splines of the Second Order of Approximation	
	$n = 10$	$n = 20$
0	0.06744487	0.06763105
0.2	0.36908780	0.36939306
0.4	0.65601636	0.65642854
0.6	0.91679162	0.91729428
0.8	1.14101729	1.14159039
1.0	1.31975420	1.32037489

Table 4. The Approximate Values of the Solution when Splines of the Second Order of Approximation were used

x_j	Splines of the Second Order of Approximation	
	$n = 40$	$n = 80$
0	0.06767764	0.06768929
0.2	0.36946943	0.36948853
0.4	0.65653166	0.65655744
0.6	0.91742003	0.91745147
0.8	1.14173376	1.14176961
1.0	1.32053017	1.32056899

Table 5. The Approximate Values of the Solution when Splines of the second order of approximation and the seventh of Approximation were used

x_j	Splines	
	Splines of the Second Order of Approximation $n = 640$	Splines of the Seventh Order of Approximation $n = 10$
0	0.067693111	0.06769317
0.2	0.36949480	0.36949490
0.4	0.65656590	0.65656604
0.6	0.91746179	0.91746195
0.8	1.14178137	1.14178156
1.0	1.32058173	1.32058194

Table 6. The Approximate Values of the Solution when Splines of the Seventh Order of Approximation were used

x_j	Splines of the Seventh Order of Approximation	
	$n = 20$	$n = 40$
0	0.06769317	0.067689317
0.2	0.36949490	0.36949490
0.4	0.65656603	0.65656604
0.6	0.91746195	0.91746195
0.8	1.14178156	1.14178156
1.0	1.32058194	1.32058194

The absolute values of the differences between the solution found on the grid of nodes ($n = 10$) and the solution found on the grid of nodes ($n = 20$) are shown in Figure 2 (blue line); absolute values of the differences between the solution found on the grid ($n = 20$) and the solution found on the grid ($n = 40$) are shown in Figure 2 (green line).




Fig. 2: The absolute values of the differences between the solution found on the grid of nodes ($n = 10$) and the solution found on the grid of nodes ($n = 20$) (blue line); absolute values of the differences between the solution found on the grid ($n = 20$) and the solution found on the grid ($n = 40$) (green line).

Figure 3 shows the absolute values of the differences of the solution found on the grid ($n = 10$) and the solution found on the grid ($n = 20$) when the splines of the seventh order of approximation were used.




Fig. 3: The absolute values of the differences of the solution found on the grid ($n = 10$) and the solution found on the grid ($n = 20$) when the splines of the seventh order of approximation were used.

Figure 4 shows the absolute values of the differences of the solution found on the grid ($n = 20$) and the solution found on the grid ($n = 40$) when the splines of the seventh order of approximation were used.

Figure 5 shows the plot of the approximate solution when the splines of the seventh order of approximation were used and $n = 40$.

The calculation results show that splines of the seventh order of approximation give an approximate solution at the grid nodes with eight correct digits in the mantissa when $n = 20$.




Fig. 4: The absolute values of the differences of the solution found on the grid ($n = 20$) and the solution found on the grid ($n = 40$) when the splines of the seventh order of approximation were used.




Fig. 5: The plot of the approximate solution when the splines of the seventh order of approximation were used ($n = 40$).

Comparing the results in the second and third columns of Table 5, we see that the spline of the second order of approximation provides five correct digits in the mantissa when we use 640 grid nodes. At the same time, to achieve the same accuracy using splines of the seventh order of approximation, it is enough to use 10 grid nodes.

4 Conclusion

In this paper, we consider the stability of the solution of the integral equations of the second kind using splines of the second and the seventh order of approximation. It should be noted that with the same number of grid nodes, polynomial splines of the seventh order of approximation provide a smaller error compared to splines of the second order of approximation. When using splines of the seventh order of approximation, a large number of nodes is not recommended.

In the future, numerical schemes for integro-differential equations will be constructed.

References:

- [1] V.R. Ibrahimov, G.Y.U. Mehdiyeva, X.G. Yue, M.K.A. Kaabar, S. Noeiaghdam, D.A. Juraev, Novel Symmetric Numerical Methods for Solving Symmetric Mathematical Problems, *International Journal of Circuits, Systems and Signal Processing*, Vol.15, 2021, pp. 1545-1557.
- [2] A. Carpio, E. Cebrian, Positivity Preserving High Order Schemes for Angiogenesis Models, *International Journal of Nonlinear Sciences and Numerical Simulation*, 2021, DOI: 10.1515/ijnsns-2021-0112.
- [3] P.Assari, F.Asadi-Mehregan, The Approximate Solution of Charged Particle Motion Equations in Oscillating Magnetic Fields using the Local Multiquadrics Collocation Method, *Engineering with Computers*, Vol.37, No. 1, 2021, pp. 21-38. DOI: 10.1007/s00366-019-00807-z.
- [4] S. Saha, V. Kumar, A. N. Das, An Elastic Half Space with a Moving Punch, *WSEAS Transactions on Applied and Theoretical Mechanics*, Vol. 16, 2021, pp. 245-249.
- [5] L. Hacıa, K. Bednarek, A. Tomczewski, Computational Results for Integral Modeling in Some Problems of Electrical Engineering, *WSEAS International Conference on Computers*, 2007, pp. 114-119.
- [6] D.T. Waide, D.G. Green, G.F. Gribakin, BSHF: A Program to Solve the Hartree–Fock Equations for Arbitrary Central Potentials using a B-spline Basis, *Computer Physics Communications*, Vol. 250, paper 107112, 2020, DOI: 10.1016/j.cpc.2019.107112.
- [7] S. Rahbar, Solving Fredholm Integral Equation Using Legendre Wavelet Functions, *WSEAS Transactions on Mathematics*, 2004, No. 3.
- [8] A. Mennouni, N.E. Ramdani, K. Zennir, A New Class of Fredholm Integral Equations of the Second Kind with Non-Symmetric Kernel: Solving by Wavelets Method, *Boletim da Sociedade Paranaense de Matematica*, Vol. 39, No 6, 2020, pp. 67-80, DOI: 10.5269/BSPM.41734
- [9] K. Maleknejad, H. Shahi Kalalagh, An Iterative Approach for Solving Nonlinear Volterra–Fredholm Integral Equation Using Tension Spline, *Iranian Journal of Science and Technology, Transaction A: Science*, Vol. 44, No 5, 2020, pp. 1531-1539, DOI: 10.1007/s40995-020-00963-8.
- [10] M.S. Islam, A. Smith, Approximating Solutions of Fredholm Integral Equations via a General Spline Maximum Entropy Method, *International Journal of Applied and Computational Mathematics*, Vol. 6, No 3, 2020, paper 64, DOI: 10.1007/s40819-020-00820-7.
- [11] A.M. Bica, C. Popescu, Fuzzy Bezier Splines with Application to Fuzzy Functional Integral Equations, *Soft Computing*, Vol.24, No 8, 2020, pp. 6069-6084. DOI: 10.1007/s00500-020-04740-y.
- [12] J. Rashidinia, K. Maleknejad, H. Jalilian, Convergence Analysis of Non-Polynomial Spline Functions for the Fredholm Integral equation, *International Journal of Computer*

- Mathematics*, Vol. 97, No 6, 2020, pp. 1197-1211.
- [13] D. Černá, V. Finěk, Galerkin Method with New Quadratic Spline Wavelets for Integral and Integro-differential Equations, *Journal of Computational and Applied Mathematics*, Vol.363, 2020, pp. 426-443. DOI: 10.1016/j.cam.2019.06.03.
- [14] M. Asif, I. Khan, N. Haider, Q. Al-Mdallal, Legendre Multi-wavelets Collocation Method for Numerical Solution of Linear and Nonlinear Integral Equations, *Alexandria Engineering Journal*, 2020, DOI: 10.1016/j.aej.2020.09.040
- [15] B. Shiri, I. Perfilieva, Z. Alijani, Classical Approximation for Fuzzy Fredholm Integral, *Fuzzy Sets and Systems*, Vol. 404, 2021, pp. 159-177.
- [16] A. Flayih Jaber, A.M.H.R. Al-Rammahi, Orthogonal Basis to Solve Linear Fredholm Integral Equations via Galerkin Method, *IOP Conference Series: Materials Science and Engineering*, Vol. 928, No 4, 2020, paper 042016, DOI: 10.1088/1757-899X/928/4/042016
- [17] M. Galina, I. Vagif, I. Mehriban, On the Construction of the Advanced Hybrid Methods and Application to Solving Volterra Integral Equation, *WSEAS Transactions on Systems and Control*, Vol. 14, 2019, pp. 183-189.
- [18] M. Galina, I. Vagif, I. Mehriban, On the Construction of the Forward-Jumping Method and its Application to Solving of the Volterra Integral Equations with Symmetric Boundaries, *WSEAS Transactions on Mathematics*, Vol. 16, 2017, pp. 295-302.
- [19] L.H. Ali, J. Sulaiman, S.R.M. Hashim, Numerical Solution of Fuzzy Fredholm Integral Equations of Second Kind Using Half-Sweep Gauss-Seidel Iteration, *Journal of Engineering Science and Technology*, Vol. 15, No 5, 2020, pp. 3303-3313.
- [20] H. Du, Z. Chen, A New Reproducing Kernel Method with Higher Convergence Order for Solving a Volterra–Fredholm Integral Equation, *Applied Mathematics Letters*, Vol. 102, 2020, paper 106117, DOI:10.1016/j.aml.2019.106117.
- [21] I.A. Dhari, N.R. Mustafa, M.M. Mustafa, Solving Linear Volterra-Fredholm Integral Equation of the Second Type Using Linear Programming Method, *Baghdad Science Journal*, Vol. 17, No 1, 2020, pp. 342-347. DOI: 10.21123/BSJ.2020.17.1(SUPPL.).0342
- [22] I.G. Burova, Fredholm integral equation and splines of the fifth order of approximation, *WSEAS Transactions on Mathematics*, Vol.21, 2022, pp. 260-270.
- [23] I.G. Burova, On Left Integro-Differential Splines and Cauchy Problem, *International Journal of Mathematical Models and Methods in Applied Sciences*, Vol. 9, 2015, pp. 683-690.
- [24] S.K. Godunov and V.S. Ryaben’kii, *Theory of difference schemes. An introduction*. North-Holland, Amsterdam; Interscience (Wiley), New York, 1964.

Contribution of Individual Authors to the Creation of a Scientific Article (Ghostwriting Policy)

-I. G. Burova developed the theoretical part,
 -G. O. Alcybeev executed the numerical experiments,
 -S. A. Schiptcova executed some numerical experiments.
 The authors are grateful to Professor V.M. Ryabov for his valuable comments.

Sources of Funding for Research Presented in a Scientific Article or Scientific Article Itself

The authors are gratefully indebted to St. Petersburg University for their financial support (Pure ID 94029567, ID 104210003)

Conflict of Interest

The authors have no conflicts of interest to declare that are relevant to the content of this article.

Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0
https://creativecommons.org/licenses/by/4.0/deed.en_US