


**NRC «Kurchatov Institute»
Saint Petersburg State University
Joint Institute for Nuclear Research**



**LXXI INTERNATIONAL CONFERENCE
«NUCLEUS-2021»**

**NUCLEAR PHYSICS AND
ELEMENTARY PARTICLE PHYSICS.
NUCLEAR PHYSICS TECHNOLOGIES.**

BOOK OF ABSTRACTS

*Online
20 – 25 September 2021*

Saint Petersburg

Organisers

NRC «Kurchatov Institute»
Saint Petersburg State University
Joint Institute for Nuclear Research

Chairs

M. Kovalchuk (Chairman, NRC “Kurchatov Institute”)

V. Zherebchevsky (Co-Chairman, SPbU)

A. Vlasnikov (Vice-Chairman, SPbU)

S. Torilov (Scientific Secretary, SPbU)

The contributions are reproduced directly from the originals.
The responsibility for misprints in the report and paper texts is
held by the authors of the reports.

International Conference “NUCLEUS – 2021. Nuclear physics and elementary particle physics. Nuclear physics technologies” (LXXI; 2021; Online part).

LXXI International conference “NUCLEUS – 2021. Nuclear physics and elementary particle physics. Nuclear physics technologies” (Saint Petersburg, Russia, 20–25 September 2021): Book of Abstracts /Ed. by V. N. Kovalenko and E. V. Andronov. – Saint Petersburg: VVM, 2021. – 396p.

ISBN: 978-5-9651-1357-6


Международная Конференция «ЯДРО – 2021. Ядерная физика и физика элементарных частиц. Ядерно-физические технологии» (LXX; 2021; Онлайн часть).

LXXI Международная Конференция «ЯДРО – 2021. Ядерная физика и физика элементарных частиц. Ядерно-физические технологии» (Санкт-Петербург, Россия, 20–25 Сентября 2021): Аннот. докл./под ред. В.Н. Коваленко, Е.В. Андропова. – Санкт-Петербург: ВВМ, 2021. – 396 с.

ISBN: 978-5-9651-1357-6

ISBN: 978-5-9651-1357-6

НИЦ «Курчатовский институт»
Санкт-Петербургский государственный университет
Объединенный институт ядерных исследований



LXXI МЕЖДУНАРОДНАЯ
КОНФЕРЕНЦИЯ
«ЯДРО-2021»

ЯДЕРНАЯ ФИЗИКА И
ФИЗИКА ЭЛЕМЕНТАРНЫХ ЧАСТИЦ.
ЯДЕРНО-ФИЗИЧЕСКИЕ ТЕХНОЛОГИИ.

СБОРНИК ТЕЗИСОВ

Онлайн
20 – 25 сентября 2021

Санкт-Петербург

INFLUENCE OF HIGH-ENERGY PROTON AND GAMMA-RADIATION ON DNA STRUCTURE IN SOLUTION

Authors: Alexandr Khalikov¹; Daria Brozhik¹; Omar Mahmoud Elsayed Kotb²; Sofia Paston³; Elena Gulevich¹; Fedor Pak¹; Jan Karlin¹; Valerij Verbenko¹; Viktor Ezhov¹

¹ Petersburg Nuclear Physics Institute of National Research Center “Kurchatov Institute”, Gatchina, Russia

² Department of Physics, Faculty of Science, Zagazig University, Egypt

³ Saint Petersburg State University, Russia

Corresponding Author: s.v.paston@spbu.ru

The use of a specific technique and type of radiation in radiotherapy of tumors is determined by various factors: the localization of the tumor in the body, its oxygenation, the stage of the disease, the availability of the technique for the patient, etc. The study of damage in the DNA structure under the influence of different types of radiation and with varying irradiation conditions provide important information for improving the methods of radiation therapy. This work compares the damage in the DNA structure under the action of ^{60}Co γ -radiation (used in the gamma knife therapeutic device) and 1 GeV protons (at synchrocyclotron of the St. Petersburg Institute of Nuclear Physics Research Center “Kurchatov Institute” SC-1000, since 1975 there has been a medical center for stereotactic proton therapy). These two types of radiation have the same value of LET = 0.3 keV/ μm .

Disturbances in the DNA structure were studied by spectrophotometric melting. The parameters of the helix-coil transition in the irradiated macromolecule are influenced by various types of radiation damage. Single- and double-strand breaks, destruction, modification, and release of nucleobases lead to destabilization of the secondary structure and lower the melting temperature of DNA (T_m), while interstrand cross-links increase T_m [1].

Melting curves of DNA irradiated with doses of 0-100 Gy were obtained at ionic strengths of solutions of 5 mM and 150 mM NaCl , as well as at a total ionic strength of 5 mM = 3 mM Na^+ + 2 mM Mg^{2+} . The absorption spectra of DNA solutions were measured at 25°C, 95°C, as well as after melting and rapid (within 10 min) cooling to 25°C. From these data, the helix-coil transition interval, T_m , the DNA molar extinction coefficient, the hyperchromic effect, and the degree of DNA renaturation after melting upon rapid cooling were determined. Proton and γ -irradiation cause a broadening of the melting interval, which indicates an increase in the heterogeneity of the DNA structure, i.e. the appearance in the DNA chain of regions that differ sufficiently in thermal stability. For DNA irradiated in 150 mM NaCl solutions with proton and γ -radiation, several maxima are observed in the differential melting curves. Under these conditions, it is possible to assume the formation of cross-links (both between two complementary strands and between DNA regions distant along the chain) as a result of irradiation. In 0.15M NaCl solutions, the secondary structure of DNA is more resistant to radiation than in 5mM NaCl solutions. It was found that proton irradiation causes a smaller drop in DNA T_m than γ -irradiation in a 0.15M NaCl solution at doses of 0–100 Gy and in a 5 mM NaCl solution at doses of 70-100 Gy. For the DNA irradiated with protons at doses of 10 and 20 Gy, an abnormally large hyperchromic effect was recorded, exceeding the value measured for native DNA. The data obtained allow us to conclude that under proton irradiation, clustered damage are formed in the DNA structure – slowly repairable or unreparable sites, while under γ -irradiation, isolated DNA lesions appear, which can be quickly and efficiently repaired in the cell [2]. Thus, irradiation with high-energy protons has a greater lethal effect on the cell than γ -irradiation.

References:

1. C.R. Cantor, P.R. Schimmel. Biophysical Chemistry. Part 2,3. W. H. Freeman and Company, San Francisco (1980).
2. E. Sage, N. Shikazono, Free Radic. Biol. Med. **107**, 125 (2017).