

Article

Algal Diversity in *Paramecium bursaria*: Species Identification, Detection of *Choricystis parasitica*, and Assessment of the Interaction Specificity

Felicitas E. Flemming ¹, Alexey Potekhin ^{2,3}, Thomas Pröschold ⁴ and Martina Schrallhammer ^{1,*}

¹ Microbiology, Institute of Biology II, Albert Ludwig University of Freiburg, 79104 Freiburg, Germany; felicitas.flemming@biologie.uni-freiburg.de

² Department of Microbiology, Faculty of Biology, Saint Petersburg State University, 199034 Saint Petersburg, Russia; alexey.potekhin@spbu.ru or loxodes@list.ru

³ Laboratory of Cellular and Molecular Protistology, Zoological Institute RAS, 199034 Saint Petersburg, Russia

⁴ Research Department for Limnology, University of Innsbruck, Mondsee, 5310 Mondsee, Austria; Thomas.Proschold@uibk.ac.at

* Correspondence: martina.schrallhammer@biologie.uni-freiburg.de

Received: 6 May 2020; Accepted: 19 July 2020; Published: 23 July 2020



Abstract: The ‘green’ ciliate *Paramecium bursaria* lives in mutualistic symbiosis with green algae belonging to the species *Chlorella variabilis* or *Micractinium conductrix*. We analysed the diversity of algal endosymbionts and their *P. bursaria* hosts in nine strains from geographically diverse origins. Therefore, their phylogenies using different molecular markers were inferred. The green paramecia belong to different syngens of *P. bursaria*. The intracellular algae were assigned to *Chl. variabilis*, *M. conductrix* or, surprisingly, *Choricystis parasitica*. This usually free-living alga co-occurs with *M. conductrix* in the host’s cytoplasm. Addressing the potential status of *Chor. parasitica* as second additional endosymbiont, we determined if it is capable of symbiosis establishment and replication within a host cell. Symbiont-free *P. bursaria* were generated by cycloheximide treatment. Those aposymbiotic *P. bursaria* were used for experimental infections to investigate the symbiosis specificity not only between *P. bursaria* and *Chor. parasitica* but including also *Chl. variabilis* and *M. conductrix*. For each algae we observed the uptake and incorporation in individual perialgal vacuoles. These host-symbiont associations are stable since more than five months. Thus, *Chor. parasitica* and *P. bursaria* can form an intimate and long-term interaction. This study provides new insights into the diversity of *P. bursaria* algal symbionts.

Keywords: *Chlorella*; endosymbiosis; intracellular algae; *Micractinium*; photobiont; infection; syngen

1. Introduction

The genus *Paramecium* Müller [1] (Peniculida, Oligohymenophorea, Ciliophora) includes two species that are known to maintain intracellular algae. *Paramecium chlorelligerum* Kahl [2,3] lives in symbiotic association with green algae belonging to the genus *Meyerella* Fawley & K. Fawley [4] (Chlorellaceae, Trebouxiophyceae, Chlorophyta) and is considered as an endemic, extremely rare species [3,5]. The other *Paramecium* species with algal symbionts is *Paramecium bursaria* Focke [6], the ‘green’ *Paramecium*, frequently found in freshwater habitats around the world [7]. Evolutionarily, this species has been assumed to be the first that diverged within the genus [8,9]. This green ciliate harbours hundreds of symbiotic green algae within its cytoplasm [7,10–14] either belonging to *Chlorella variabilis* Shihira & R. W. Krauss [15] or to *Micractinium conductrix* (K. Brandt) Pröschold & Darienko [16,17] (Chlorellaceae, Trebouxiophyceae, Chlorophyta).

The algae escape lysosomal fusion by incorporation in perialgal vacuoles [18–20]. In this facultative, mutualistic symbiosis, the algae provide photosynthesis products for their host [21–23] and also form a protective layer against UV-radiation damage [24]. Moreover, they profit from protection against the lytic *Paramecium bursaria* chlorovirus [25,26]. In addition, the host increases its symbiont's motility by providing transport to brightly illuminated areas optimal for photosynthesis [27]. The facultative nature of this symbiosis allows the separate cultivation of both organisms, making the association an easy-to-access model system to study symbiotic interactions [28–30]. Albeit under laboratory conditions the symbiosis is not obligate, it is the natural condition in the environment. Both algae have not been retrieved as free-living organisms from nature so far and, therefore, are considered as highly adapted to their endosymbiotic lifestyle [31]. Similarly, aposymbiotic *P. bursaria* cells have been isolated from natural freshwater sources only on rare occasions [32]. As *Chl. variabilis* and *M. conductrix* apparently are mainly restricted to their endosymbiotic lifestyle within *P. bursaria*, these associations have been considered as highly specific. However, recent studies report these microalgae as intracellular symbionts from different host organisms such as *Tetrahymena* [33], *Hydra* [34], *Frontonia vernalis* [35], and sponges [10]. Furthermore, *P. bursaria* can also harbour different organisms either co-occurring with *Chlorella*-like algae, i.e., bacteria [36,37], or in their place, i.e., yeasts [38,39]. Therefore, the symbiotic relationship between *P. bursaria* and its intracellular green algae might not be as exclusive as previously assumed.

An interesting case of double algal infection in *P. bursaria* by *Chlorella* and another green alga was reported by two Japanese groups [31,40]. The intracellularly uncommon, additional alga was characterized as *Chlorocystis parasitica* (K. Brandt) Pröschold & Darienko [16,17] (Trebouxiophyceae, Chlorophyta). Cells of this alga were observed between the trichocysts near the host's cortex [31,40]. This localization corresponds to that of the natural algal endosymbionts'. The cortex area is devoid of lysosomes, thus an intracellular symbiont is likely protected here against lysosomal fusion [41] and hence against host defence mechanisms, which was interpreted as an advanced symbiotic status [31,40]. Due to its small cell size (1.5–3.0 μm in length, 1.0–1.5 μm in width; [10]), *Chor. parasitica* will be referred to as picoalga.

Characterization of symbiotic systems often requires interdisciplinary expertise to fulfill the standards of the respective scientific disciplines. Molecular characterization of a single or more phylogenetic markers is typically part of current descriptions of new taxa. Sequence analysis allows evolutionary interpretation in absence of fossil records and furthermore facilitates the assignment of new isolates to described taxa. The choice of the molecular markers depends on multiple factors and varies between organism groups and the desired level of phylogenetic resolution. In case of *P. bursaria*, the SSU (small subunit or 18S ribosomal RNA) gene sequence is commonly applied for phylogenetic analyses of members of the genus *Paramecium* [5,42–44]. Some recognized *Paramecium* species are actually sibling or cryptic species complexes, i.e., their members are morphologically indistinguishable but reproductively isolated and thus are called syngens. The most prominent example are the syngens of the *Paramecium aurelia* complex, which was first studied in detail by Sonneborn [45]. Those syngens are nowadays recognized as separate species. Similarly, the existence of multiple syngens was described for *P. bursaria* [46]. Bomford's description of *P. bursaria* has been the most extensive one, unfortunately his strain collection was lost. Only few strains remain and are scattered across different laboratories. A representative collection of *P. bursaria* strains assigned to five syngens, R1 to R5 [47], is maintained at the RC CCM collection (World Data Centre for Microorganisms, RN 1171), Saint Petersburg State University, Saint Petersburg, Russia. Molecular phylogenetic analyses confirmed *P. bursaria* to be a complex of at least five cryptic species [47–50] using different molecular markers (e.g., ITS1-5.8S-ITS2-5'LSU fragments, mitochondrial cytochrome c oxidase subunit I gene, histone H4 gene; [5,48,51]).

Based on molecular phylogenetic analyses of SSU and ITS (internal transcribed spacer) regions of nuclear-encoded rRNA genes, the genus and species concepts within the Chlorellaceae remain provisional due to the lack of bootstrap support in molecular analyses [10,52–55]. The family is divided

into two major clades: (i) the well-supported *Parachlorella* clade and (ii) the moderately supported *Chlorella* clade [56,57]. Furthermore, the ITS2 region has been used for species delineation of members of the Chlorellaceae [10,50,55,58]. The two common endosymbionts of *P. bursaria*, *Chl. variabilis* and *M. conductrix*, both vary in the ITS2 secondary structure and can be differentiated using their ITS2 barcodes [50].

This study addresses the diversity of intracellular algae in *P. bursaria*. Therefore, ciliate hosts and algal endosymbionts were identified using different molecular markers. In order to analyse the interaction between the three different algal species detected and *P. bursaria*, experimental re- and cross-infections were performed and the fate of the intracellular algae was monitored. Differences in the establishment of stable associations will allow to draw conclusions regarding the compatibility of the different algae as endosymbionts of *P. bursaria* and the specificity of such interactions.

2. Materials and Methods

2.1. Strains and Cultivation

All *P. bursaria* strains (Table 1) were cultivated in Dryl's solution (0.1 M Na₂HPO₄, 0.1 M NaH₂PO₄, 0.1 M CaCl₂, 0.1 M sodium citrate, pH 6.8) at 20 °C in a light chamber (EHRET Type KLT/S 4, Ehret, Emmendingen, Germany) with a 12 h/12 h light-dark cycle. The cultures were fed weekly with the bacterium *Raoultella planticola* DMSZ 3069 as food organism (bacterized CM). Therefore, the bacteria were inoculated in 0.25% Cerophyll medium (CM; [59] prepared from wheat grass powder (GSE-Vertrieb), Saarbrücken, Germany) and a stigmasterol (Sigma-Aldrich, Munich, Germany) concentration of 500 µg/L. After two days of incubation at 20 °C, the bacterized CM was ready for use.

Table 1. *Paramecium bursaria* and algal strains used in this study. Obtained sequences and their accession numbers are indicated.

Strain	Host	Syngen	Acc. Number (Region)	Algal Symbiont	Acc. Number (Region)	Collection Date	Collected by
JPN ¹	<i>P. bursaria</i>	R3	MT460147 (SSU-ITS-LSU part.)	<i>Chlorella variabilis</i>	MT460236 (SSU-ITS)	September 2014	L. Koehler
Tue2015 ²	<i>P. bursaria</i>	R3	MT460146 (SSU-ITS-LSU part.)	<i>Chlorella variabilis</i>	MT460235 (SSU-ITS)	-	K. Eisler
Frieds ³	<i>P. bursaria</i>	R2	MT460149 (SSU-ITS-LSU part.)	1st <i>Micractinium conductrix</i>	MT460238 (SSU-ITS)	August 2014	M. Witt
				2nd <i>Choricystis parasitica</i>	MT459641 (partial SSU)		
Old-Pf ⁴	<i>P. bursaria</i>	R1	MT459459 (SSU-ITS-LSU part.)	<i>Micractinium conductrix</i>	MT460234 (SSU-ITS)	August 2014	K. Grosser
RanNy ⁵	<i>P. bursaria</i>	R2	MT460150 (SSU-ITS-LSU part.)	<i>Micractinium conductrix</i>	- (diagn. PCR)	Jul. 2016	K. Grosser
Scot ⁶	<i>P. bursaria</i>	R1	MT460148 (SSU-ITS-LSU part.)	<i>Micractinium conductrix</i>	MT460237 (SSU-ITS)	July 2010	M. Schrallhammer
Ek ⁷	<i>P. bursaria</i>	R2	MT576702 (SSU-ITS-LSU part.)	<i>Micractinium conductrix</i>	- (diagn. PCR)	September 2009	A. Potekhin
Bob2 ⁸	<i>P. bursaria</i>	R2	MT576703 (SSU-ITS-LSU part.)	<i>Micractinium conductrix</i>	- (diagn. PCR)	August 2006	A. Potekhin
Ard10 ⁹	<i>P. bursaria</i>	R4	MT576704 (ITS-LSU part.)	<i>Chlorella * variabilis</i>	KM203667 (LSU partial)	April 2006	V. Yashchenko

¹: Pond at Rurikojo temple, Yamaguchi, Japan; ²: Culture collection of Dr. K. Eisler, University of Tübingen, Germany; ³: Friedrichsee, Saxony-Anhalt, Germany; ⁴: Pond 'Pferdetränke', Oldenburg, Germany; ⁵: Nymphensee, Rangsdorf, Germany; ⁶: Loch Katrine, Scotland, United Kingdom; ⁷: Saint Petersburg, pond in the park, Russia; ⁸: Forest pond, Vyborg, Saint Petersburg region, Russia; ⁹: Pond, Ardmore, Oklahoma, USA; * Zagata et al., 2016 [60]; part.: partial.

2.2. Observation and Size Determination of Intracellular Algae

The intracellular algae were investigated by detecting the autofluorescence of the chlorophyll by fluorescence microscopy (Axio Imager.M2, Zeiss, Jena, Germany). Therefore, *Paramecium* cells were

fixed with Bouin's fixative solution (Sigma-Aldrich) and transferred to a microscope slide. Cells were examined using the Cy5 channel (EX BP 640/30, BS FT 660, EM BP 690/50) and an exposure time ranging from 3000 to 7000 ms in order to discriminate between the individual intracellular algae. To determine the size of the intracellular micro- and picoalgae, the diameter of 15 cells per strain was measured.

2.3. DNA Extraction and Amplification of Molecular Markers

All polymerase chain reactions (PCR) were performed with a PeqStar 2× thermocycler (Peqlab, VWR International, Darmstadt, Germany) using TaKaRa reagents and TaKaRa ExTaq polymerase (TaKaRa Bio, Otsu, Japan). Primer sequences (Table 2) and amplification protocols (Supplementary Tables S1 and S2) are provided. The SSU-ITS1-5.8S-ITS2-5'LSU rRNA gene region of *P. bursaria* was amplified using the primer combinations Penic_F82 and Penic_R1280 and Penic_F661 and 28S_R457. Primers Penic_F82, Penic_F661 and 28S_R457 were used for sequencing.

Table 2. Oligonucleotides used for molecular characterization of both *P. bursaria* and symbiotic algae.

Primer	Sequence [5'-3']	Literature
Penic_F82	GAAACTGCGAATGGCTC	Strüder-Kypke et al., 2000 [8]
Penic_F661	ATAGATGGGGGCATTAGT	mod. from Fokin et al., 2006 [61]
Penic_R1280	CGACACGTCCTAACAGA	Fokin et al., 2006 [61]
28S_R457	CTTCCTTCGYAGTACT	W. Ludwig, pers. commun.
AF	TCGACAATCTGGTGATCCTGCCAGT	Pröschold et al., 2001 [10]
Chlo_F238	GCCCTATCAACTTTCGATG	this study
Chlo_G500F	GAATGAGTACAATCTAAACCCCTTAAC	Darienko et al., 2019 [62]
Chlo_G800F	CCTGTTGGTCTGTAGGAGTGGAGTAATG	Darienko et al., 2019 [62]
Chlo_F1074	GGGTTGCCTTGTCAAGG	this study
ITS055R	CTCCTTGGTCCGTGTTCAAGACGGG	Marin et al., 2003 [63]
Chlo_G800R	CATTACTCCGCTCCTACAGACCAACAGG	Darienko et al., 2019 [62]
Chlo_R841	CGGAGTCATCGAAGAAC	this study
Chori_F238	GCCCTATCAACTTCAACC	this study
Chori_R841	TGGGGGGGTCATCAAAGG	this study

For the intracellular algae, the nuclear encoded SSU gene and the ITS region were amplified using a semi-nested PCR approach to obtain sufficient DNA amounts for direct sequencing. For the initial amplification (primer combination: AF and ITS055R) as well as for the two semi-nested amplification reactions (primer combinations: AF and Chlo_R841 and Chlo_G800F and ITS055R) the protocol described by Pröschold and colleagues [10] was used. AF, Chlo_F238, Chlo_G800F, Chlo_R841 and ITS055R were used as sequencing primers. For strains Scot and Old-Pf amplification was carried out with primer combination Chlo_G800F and ITS055R (Supplementary Tables S1 and S2) and both primers for sequencing. Additionally, a diagnostic PCR allowing to discriminate between *Chl. variabilis* and *M. conductrix* was carried out as described elsewhere [50].

To amplify the SSU gene of *Choricystis* from strain Frieds the primer combination Chori_F238 and Chori_R841 was used. Those primers were designed to specifically match *Choricystis*-like algae based on preliminary sequence data. Chori_F238 was used for sequencing.

Purified PCR products (QIAquick PCR Purification Kit, Qiagen, Hilden, Germany) were sequenced at GATC Biotech (Konstanz, Germany). Accession numbers of obtained sequences are provided in Table 1.

2.4. Molecular Characterization of *Paramecium bursaria*

In order to confirm our morphological identification of the ciliates as members of *P. bursaria*, we sequenced the SSU rRNA gene sequence and the ITS region of our newly obtained strains. Both markers are commonly applied for characterization at generic and species level [5,42–44,47,50].

The obtained sequences were imported into two datasets: (i) The SSU sequences were included into a dataset of representatives of most *Paramecium* species received from the SILVA SSU Ref NR 99 release 138 database [64]. In addition, nine other members of Peniculidae were included as outgroup resulting in a dataset of 59 sequences with a length of 1493 bp. (ii) The ITS sequences were included into an alignment of 32 *P. bursaria* strains (506 bp) as described in previous studies [47,50]. For syngen assignment, we added sequences of additional strains (Ard10, Bob2, Ek), which were tested for their mating behaviour by RC CCM to increase the number of sequenced strains per syngen. The ITS sequences were aligned according to their secondary structures.

The best fitting evolutionary models for these datasets were determined using the automated model selection tool implemented in PAUP (version 4.0a, built 167) [65]. The best model (SSU: GTR+I+G; ITS: K81uf+I) was chosen following the Akaike Information Criterion (AIC). For the Bayesian analyses, Bayesian Information Criteria (BIC) were chosen (SSU: GTR+I+G; ITS: GTR+I) for calculation in MrBayes (version 3.2.6 x64; [66]). Non-parametric Maximum Likelihood (ML) analysis was estimated on 1000 pseudoreplicates (PHYML 2.4.5 from the ARB software package; [67]). Bayesian Interference (BI) analysis was carried out running three runs with one cold and three heated Markov Chain Monte Carlo chains for 1,000,000 generations and sampling the first 25% of the generations as burn-in.

2.5. Molecular Characterization of Green Algal Endosymbionts in *Paramecium bursaria*

For the molecular characterization of the algal endosymbionts we used sequences covering the SSU-ITS1-5.8S-ITS2 region for discrimination at species level in accordance with previous studies [10,55,56,62,68,69]. Especially the ITS2 region of this nuclear encoded ribosomal operon is often used for species discrimination between members of the Chlorellaceae [53,55,56,70]. The here performed amplification of the ITS2 region covered helices I-III but only partial helix IV. The analysed dataset comprised 50 sequences of representative members of the Chlorellaceae, which were aligned according to the secondary structures predicted by the software mfold [71], following the approach of Pröschold and colleagues [10,55]. This software used the thermodynamic model (minimal energy) for RNA folding. The best fitting evolutionary model for this dataset was predicted and ML and BI analyses were conducted as described above except that BI analysis was run for 5,000,000 generations. Additionally, the Neighbour Joining method (implemented in PAUP) was used. The analysis was run with 1000 bootstrap replicates and a 50% majority rule consensus tree was calculated.

Furthermore, we compared the algal endosymbionts' secondary structures of ITS2 helices I-III to that of the type strains of *Chl. variabilis* and *M. conductrix*. In case of the symbionts of strains Bob2, Ek, and RanNy these algae were identified by means of diagnostic PCR (Supplementary Figure S1).

To identify the intracellular picoalgae of *P. bursaria* observed in strain Frieds, we used the obtained partial SSU gene sequence and compared it with 41 representative sequences for the *Trebouxia* lineage (Trebouxiophyceae; following Darienko and colleagues [62]) with the *Prasiola* clade (four species) and the genus *Neocystis* (two species) as outgroup. The alignment was obtained as described for *Paramecium* SSU and comprised 948 positions. Model selection (TIM+I+G) for this dataset was performed as detailed above. Since this model is not implemented in MrBayes, it was substituted with GTR+I+G. ML and BI analyses were performed as described above for *P. bursaria*. The settings of all used evolutionary models are provided in Supplementary Table S3.

2.6. Establishment of Aposymbiotic *Paramecium bursaria*

Algal-free *P. bursaria* were generated by treatment with cycloheximid (Roth, Karlsruhe, Germany). Therefore, symbiotic *P. bursaria* cultures not fed for at least seven days were incubated in 10 µg/mL cycloheximid solution and kept under constant light conditions for approximately three days, then the cells were washed with Dryl's solution by filtering over a 10 µm membrane and fed with bacterized CM. This filtering step was repeated three to five times over the course of two weeks to assure complete elimination of the algae. If necessary, a second consecutive cycloheximid treatment was performed seven to ten days after the initial exposure. Microscopic inspection via fluorescence microscopy

confirmed the elimination of intracellular algae. Aposymbiotic status was confirmed if after 10 seconds of exposure time no autofluorescence signal of the algae's chloroplasts was detected.

2.7. Pulse-Chase Infection Experiments

To investigate the symbiosis specificity between *P. bursaria* and three different symbiotic algal species, we exposed aposymbiotic cells (strains JPN, RanNy, Scot; affiliated to different syngens, see below) to isolated *Chl. variabilis*, *M. conductrix*, or *Chor. parasitica* (Figure 1). Aposymbiotic *P. bursaria* generated by treatment with cycloheximide were exposed (recipient) to isolated algae obtained from symbiotic *P. bursaria* cultures (donor; Supplementary Table S4). In re-infection assays, the supplied algae belonged to the same algal species as the recipient strain originally harboured, while in cross-infection experiments a different species was provided. We followed the fate of the algae after their uptake and differentiated between digestion, expulsion, and endosymbiotic maintenance by fluorescence microscopy using the above listed parameters. Each experiment comprised three replicates. In total, 20 aposymbiotic *Paramecium* cells were exposed to a suspension of the respective alga for 5 min with a ratio of 3×10^3 algae per ciliate cell. Then the paramecia were washed and transferred to 200 μL Dryl's solution inoculated with 20 μL of bacterized CM. 20 μL bacterized CM were added every second day for seven to ten days. Afterwards, the amount of supplied food was adjusted with respect to cell densities. A regular weekly feeding schedule was implemented after two to three weeks post infection (p.i.). The establishment of the symbiosis was monitored using the fluorescence signal of the algae's chlorophyll as described above. Instead of Bouin's solution, the *Paramecium* cells were fixed with 2% paraformaldehyde (PFA), which resulted in lower background signal and thus an improved signal to noise ratio. To monitor the process of re-establishment of symbiosis, *P. bursaria* cells were microscopically screened after two, four, seven, ten, and 14 days. Successful infection was confirmed when individually enclosed algae were localized beneath the host's cell cortex.




Figure 1. Pulse-chase infection experiments to assess the symbiosis specificity between *Paramecium bursaria* and its symbiotic algae. Symbiotic cultures of *P. bursaria* (donor) were mechanically lysed to obtain algal suspensions of *Chlorella variabilis* and *Micractinium conductrix*, which were used in re- and cross-infection experiments, respectively. Detailed information about strains used in the experiments (algae as well as host) is provided in Supplementary Table S4. Algal culture of *Choricystis parasitica* was obtained via mechanical lysis of *P. bursaria* strain Frieds for subsequent extracellular cultivation. This algal culture was then used in infection experiments with aposymbiotic *P. bursaria* cells (recipient). After 5 min of exposure (pulse), the paramecia were washed to remove extracellular algae and placed into fresh medium inoculated with food bacteria (chase). Status of symbiosis was monitored regularly via autofluorescence microscopy.

3. Results

3.1. Microscopic Investigation of the Intracellular Algae

Microscopic observations of the nine *P. bursaria* strains (Table 1) confirmed the presence of intracellular green algae (data not shown). In case of strain Frieds, algal cells belonging to two size categories were detected (Figure 2). Algae belonging to the larger category showed cell diameters of $4.08 \pm 0.02 \mu\text{m}$ (referred to as microalgae) while the smaller algae had a diameter of $1.40 \pm 0.20 \mu\text{m}$ (picoalgae). The picoalgae were localized individually throughout the cytoplasm, especially near the host's cell cortex. In addition, several of these picoalgae were observed aggregated in digestive vacuoles (Figure 2B). In a single occasion we even followed the fate of a digestive vacuole comprising more than 20 algal cells being emptied into the surrounding medium and thus *Chor. parasitica* were expelled.




Figure 2. *Paramecium bursaria* and its intracellular algae. (A) *P. bursaria* cell of strain Frieds with micro- and picoalgae distributed throughout the cytoplasm. (B) Digestive vacuole of the same cell incorporating multiple picoalgae. (C) Micro- and picoalgae simultaneously situated within the host's cytoplasm. Arrows indicate intracellular picoalgae. (D) Autofluorescence signal of the chloroplast from intracellular picoalgae of *P. bursaria* strain Frieds. Scale bars: 20 μm .

3.2. *Paramecium bursaria* and the Five Syngens

The phylogenetic inference of *Paramecium* species based on SSU rRNA gene sequences (Figure 3) recovers the expected topology and composition of the five subgenera of this genus. All here obtained sequences cluster within the monophyletic and maximal supported subgenus *Chloroparamecium*. Subgroups within the *P. bursaria* clade can be observed (Figure 3), but the achieved resolution does not allow unambiguous identification of syngen affiliations. In order to discriminate between the five syngens of *P. bursaria*, we analysed the ITS1-5.8S-ITS2-5'LSU region alone (Figure 4). While the here characterized sequences span from SSU till the beginning of the LSU (except for strain Ard10), the majority of publicly available sequences of *P. bursaria* strains with known sygen affiliation are limited to the ITS region. Sequences obtained here cluster either with (Figure 4) *P. bursaria* sygen R1 (Old-Pf and Scot), R2 (Bob2, Ek, Frieds, and RanNy), R3 (JPN and Tue2015), or R4 (Ard10). A potential correlation between the host's sygen affiliation and the present algal symbiont can be observed. Strains affiliated to syngens R3, R4, and R5 harbour *Chl. variabilis* while those assigned to sygen R2 contain *M. conductrix*. Only in case of sygen R1 both algae have been described as symbionts.




Figure 3. Molecular phylogeny of the genus *Paramecium* based on SSU rRNA gene sequences. Maximum likelihood tree based on 1493 aligned positions. GTR+I+G was used as evolutionary model. The respective subgenera are highlighted. Other members of Peniculida were used as outgroup. Bootstrap values above 70% and Bayesian Interference values above 0.95 are indicated. Asterisks indicate maximum support in both analyses. Numbers in brackets represent the number of sequences included in collapsed groups. Sequences marked in bold were obtained in this study.




Figure 4. Sygen affiliation of *Paramecium bursaria* strains based on the internal transcribed spacer (ITS) region spanning ITS1-5.8S-ITS2. Unrooted Maximum likelihood tree based on 506 aligned positions is shown. GTR+I was used as evolutionary model. Bootstrap values above 70 % and Bayesian Inference values above 0.95 are indicated. The five syngens are highlighted as R1 to R5, a plus marks a strain whose mating behaviour was previously experimentally determined. The identity of the algal symbiont if known is provided. Sequences marked in bold were obtained in this study.

3.3. *Chlorella variabilis*, *Micractinium conductrix*, and *Choricystis parasitica* as *Paramecium*'s Endosymbionts

Phylogenetic inference of the intracellular microalgae of *P. bursaria* based on sequences spanning the SSU-ITS1-5.8S-ITS2 region (Figure 5) reveals the genera *Chlorella* and *Micractinium* as two distinct monophyletic clades. The intracellular microalgal strains JPN and Tue2015 cluster with other *Chl. variabilis* strains with maximum support in the phylogenetic analyses. Frieds, Old-Pf, and Scot affiliate with sequences belonging to *M. conductrix* with maximum support.

Comparing the conserved regions of ITS2 helices I-III of *Chl. variabilis* (barcode CVAR, [50]) to that of strains JPN and Tue2015, respectively *M. conductrix* (barcode MCON, [50]) to strains Frieds, Old-Pf, and Scot, no differences were observed (data not shown).

The sequence of the second intracellular alga detected in strain Frieds clusters with other sequences of the genus *Choricystis* as monophyletic sister group to the *Elliptochloris* clade with high support in both analyses (Figure 6). The obtained sequence is identical to that of several other *Chor. parasitica* strains.




Figure 5. Molecular identification of microalgal endosymbionts. Phylogeny of members of the Chlorellaceae based on sequences covering the SSU rRNA gene and the internal transcribed spacer region spanning ITS1-5.8S-ITS2. The shown tree was inferred by the Neighbour Joining method based on 2224 aligned positions. GTR+I+G was used as evolutionary model. Bootstrap values above 70 % and Bayesian Interference values above 0.95 are indicated (ML/BI/NJ). Asterisks indicate maximum support in all analyses. Members of the *Parachlorella* clade were selected as outgroup. Black circles indicate species capable of living in symbiosis. The sequences marked in bold were obtained in this study.

3.4. Establishment of Symbioses

We followed the fate of isolated algae after their uptake and differentiated between digestion, expulsion, and endosymbiotic maintenance by aposymbiotic *P. bursaria* cells via fluorescence microscopy. Within five days after exposure of aposymbiotic *P. bursaria* cells (Figure 7A,D), digestive vacuoles were observed in all paramecia predominantly occurring in the central part of the *Paramecium* cell with more than one enclosed alga regardless of the supplied species. Multiple perialgal vacuoles enclosing single algal cells were observed throughout the entire host cytoplasm. After 12 to 15 days p.i., numerous perialgal vacuoles were localized in the cytoplasm near the host's cell cortex in each examined *Paramecium* cell (Figure 7B,E). Successful re-establishment of symbiosis was achieved for

all tested combinations (Supplementary Table S4) regardless of host background (former *Chlorella* or *Micractinium* host), syngen affiliation, and algal identity (*Chl. variabilis*, *M. conductrix*, or *Chor. parasitica*). Thus, we observed no differences between re- versus cross-infection and no preference of certain *P. bursaria* syngens for specific algae. After four months of cultivation, infected *P. bursaria* cells harbour numerous (200–500) intracellular algal symbionts. The observed symbiotic conditions are comparable to the natural state (Figure 7C,F). These cultures are stable since over five months and are maintained to date in the laboratory.




Figure 6. Molecular phylogeny of *Choricystis parasitica* and closely related members of the Chlorellaceae based on partial SSU gene sequences. Maximum likelihood tree based on 948 aligned positions. GTR+I+G was used as evolutionary model. Bootstrap values above 70 % and Bayesian Inference values above 0.95 are indicated. Asterisks indicate maximum support in both analyses. Dark circles indicate species capable of living in symbiosis, in case of *Choricystis* the symbiotic strains are highlighted. Species belonging to the *Prasiola* and *Neocystis* clades were chosen as outgroup. The obtained sequence is marked in bold.




Figure 7. Establishment of symbiosis between aposymbiotic *Paramecium bursaria* and intracellular algae. Re-establishment of symbiosis between aposymbiotic *P. bursaria* RanNy with *Micractinium conductrix* obtained from strain Scot; (A–C) and aposymbiotic *P. bursaria* JPN with *Chorocystis parasitica* (D–F). Aposymbiotic *Paramecium* cells (A,D) obtained after treatment with cycloheximide. The white outline corresponds to the ciliate’s cell shape based on bright field microscopy (not shown). Symbiotic status of *P. bursaria* after four to seven days (B,E). Successfully established symbiosis after two (C) to four (F) weeks. Representative examples are shown. Arrows indicate digestive vacuoles. Scale bars: 20 μ m.

4. Discussion

4.1. Symbiotic Relationships and Specificity of *Paramecium bursaria* and Its Green Algal Endosymbionts

Previous studies showed that aposymbiotic *P. bursaria* are able to re-establish symbiosis with their intracellular microalgae [19,72]. After uptake via phagocytosis, most of the algae are incorporated in digestive vacuoles. Some manage to escape the host’s lysosomal fusion by ‘budding off’ into the cytoplasm being enclosed in perialgal vacuoles. Subsequently, those horizontally transmitted algae are distributed throughout the host cell as well as beneath the cell cortex [73]. Based on the characterization of natural *P. bursaria*-alga symbioses, it was speculated that the occurrence of certain intracellular alga species might depend on the geographic origin of *P. bursaria*. The observed endosymbionts separated into two groups [10,12,13,74–76], i.e., the “American/Southern” (later on described as *Chl. variabilis*) and the “European/Northern” (*M. conductrix*) group. Thus, a correlation between endosymbiont and host population was assumed. Another recent study presents contradicting results reporting the first *Chl. variabilis* from European *P. bursaria* [50].

Actual testing the specificity of symbiotic partners accepted by *P. bursaria* ([77], this study) revealed that all tested intracellular algae derived from this ciliate were accepted as endosymbionts. On the other hand, free-living algae and those obtained from *Hydra* were digested [77]. A competition experiment revealed a clear preference of the aposymbiotic *P. bursaria* strain for its native *Chlorella* strain [77]. In the here conducted re- and cross-infection experiments without of a competitive setting, algae-free *P. bursaria* did not show a preference for their original symbiont or symbiont species. Aposymbiotic strains formerly harbouring *Chl. variabilis* maintained *M. conductrix* as endosymbiont, and vice versa. The picoalga *Chor. parasitica*, naturally sharing the cytoplasm of its host with *M. conductrix*, was able to form a stable symbiosis independently of co-occurring microalgae and with cells naturally harbouring *Chl. variabilis*. Even though *P. bursaria* establishes an intimate and long-term stable symbiotic association with *Chor. parasitica*, the role of the picoalga for *P. bursaria* remains unclear. An evaluation of

the symbiotic relationship between *P. bursaria* and the intracellular picoalga *Chor. parasitica* awaits further detailed analyses as current findings do not allow a placement of this interaction on the mutualism-parasitism continuum.

4.2. Molecular Characterization of *Paramecium bursaria* and Its Green Algal Endosymbionts

Unsurprisingly, the novel isolates of ‘green’ ciliates were unambiguously confirmed as *P. bursaria*. The phylogenetic tree inferred from SSU sequences is in overall good agreement with those previously published [5,42,44]. It already indicates intraspecific diversity, which becomes obvious in the analysis of the ITS region whose results reflect previous findings [47,50]. The analysis of the currently available ITS data might be interpreted that certain *P. bursaria* syngens harbour preferentially certain algal symbiont species. On the other hand, the number of symbiotic systems with both partners fully characterized is small, therefore this result should be considered very cautiously. It is likely that the picture will change with an increased amount of characterized strains. A revision of the *P. bursaria* species complex and conclusions about the suitability of certain syngens as hosts for specific algae await future studies.

The here characterized algal symbionts *Chl. variabilis* and *M. conductrix* are morphologically nearly identical [50]. However, they clearly differ in size from the additional algal symbiont detected in strain Frieds. The observed sizes in case of the microalgae were at the minimal range described for *M. conductrix* (5.0–12.0 µm, [10]) and in the typical size range for intracellular *Chor. parasitica* (1.5–3.0 µm) in case of the picoalgae. In the phylogenetic inference based on SSU-ITS sequences, the genera *Chlorella* and *Micractinium* were recovered as monophyletic and maximum support was obtained for the associations of the here studied microalgae with either *Chl. variabilis* or *M. conductrix*. The tree topology is nearly identical to that of previous studies [62,68] except for small differences lacking support in each analysis. The recovery of both algal species as endosymbiont of *P. bursaria* was not surprising, the detection of picoalgae in strain Frieds and their identification as *Chor. parasitica* more so. This is the first report of these picoalgae co-occurring with *M. conductrix* in the cytoplasm of *P. bursaria* and only the third report of *Chor. parasitica* as potential endosymbiont of *P. bursaria*. In previous observations the co-occurring primary symbiont was identified as *Chlorella* [31,40].

5. Conclusions

Paramecium bursaria is well adapted to harbour green algae as symbionts. Almost exclusively, either *Chl. variabilis* or *M. conductrix* are found in high abundances in the cytoplasm of this ciliate. One route to address the specificity of these associations is the molecular characterizations of host and symbionts. This approach provides information which symbioses are ecologically and evolutionary successful in the studied habitat. It is important to stress that therefore host and symbiont should be characterized in parallel, which is not always the case. Molecular approaches need to be combined with microscopical observations to ensure that indeed all partners are accounted for and none is overlooked or dismissed as environmental contamination. This might easily happen with minor symbionts in multiple infections or with much smaller and intracellularly less abundant picoalgae. Experiments to test the ability for symbiosis formation provide an additional perspective to the observed occurrence of associations in the environment. Infection assays can examine which interactions can be formed at e.g., species or genotype level. Thus, they provide insights into the genetic broadness of potentially realizable symbioses under constant laboratory conditions and allow to entangle the impact of additional biotic or abiotic factors shaping the formation and occurrence of symbioses. The present work can serve as a roadmap how such analyses can be conducted in regard to molecular as well as physiological characterization.

Supplementary Materials: The following are available online at <http://www.mdpi.com/1424-2818/12/8/287/s1>, Table S1: Primer combinations and their PCR program specifications. Table S2: General PCR program to amplify the SSU rRNA gene. Table S3: Model parameters as defined by PAUP. Table S4: Combinations of aposymbiotic

P. bursaria strains and algae used in the re- and cross-infection experiments in this study. Figure S1: Diagnostic PCR for identification of *Micractinium conductrix*.

Author Contributions: Conceptualization and supervision, M.S.; methodology, F.E.F., T.P., and M.S.; formal analysis, F.E.F.; resources, A.P., T.P., and M.S.; data curation, F.E.F. and M.S.; visualization, F.E.F.; funding acquisition, F.E.F., T.P., and M.S.; project administration, T.P. and M.S.; writing—original draft preparation, F.E.F. and M.S.; writing—review and editing, F.E.F. and M.S. All authors read and approved the final version of the manuscript.

Funding: This research received external funding from the Albert Ludwig University of Freiburg (Innovationsfond, project number 2100297401; MS), the Austrian Science Fund (FWF, project P28333-B25; TP), the Wissenschaftliche Gesellschaft Freiburg i. Breisgau (MS), the Grünwald-Zuberbier-Stiftung (FEF), and from the European Community's H2020 Programme H2020-MSCA-RISE 2019 under grant agreement number 872767 (MS).

Acknowledgments: We thank Bettina Sonntag for the opportunity to work at the Limnological Research Station Mondsee, Austria. We also thank Tatyana Darienko for her kind assistance with the morphology of green algae and in addition Birgit Süßenbach for preliminary molecular work and Chantal Bleile for her contributions to graphical design.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Müller, O.F. *Vernium Terrestrium et Fluvialium, seu Animalium Infusoriorum, Helminthicorum et Testaceorum non Marinorum Succincta Historia*; Heineck et Faber: Havniae et Lipsiae, Germany, 1773; Volume 1, p. 74.
2. Kahl, A. Urtiere oder Protozoa. I. Wimpertiere oder Ciliata (Infusoria). 4. Peritricha und Chonotricha. *Die Tierwelt der angrenzenden Meeresteile* **1935**, 30, 651–886.
3. Kreutz, M.; Stoeck, T.; Foissner, W. Morphological and molecular characterization of *Paramecium (Viridoparamecium) nov. subgen.) chlorelligerum* Kahl (Ciliophora). *J. Eukaryot. Microbiol.* **2012**, 59, 548–563. [[CrossRef](#)] [[PubMed](#)]
4. Fawley, M.W.; Fawley, K.P.; Owen, H.A. Diversity and ecology of small coccoid green algae from Lake Itasca, Minnesota, USA, including *Meyerella planktonica*, gen. et sp. nov. *Phycologia* **2005**, 44, 35–48. [[CrossRef](#)]
5. Lanzoni, O.; Fokin, S.I.; Lebedeva, N.; Migunova, A.; Petroni, G.; Potekhin, A. Rare freshwater ciliate *Paramecium chlorelligerum* Kahl, 1935 and its macronuclear symbiotic bacterium “*Candidatus Holospora parva*”. *PLoS ONE* **2016**, 11, e0167928. [[CrossRef](#)]
6. Focke, G. Ueber einige Organisationsverhältnisse bei polygastrischen Infusorien und Räderthieren. *Oken. Isis* **1836**, 10, 785–787.
7. Fujishima, M. *Endosymbionts in Paramecium*; Springer Science and Business Media: Berlin, Germany, 2009; Volume 12.
8. Strüder-Kypke, M.C.; Wright, A.D.G.; Fokin, S.I.; Lynn, D.H. Phylogenetic relationships of the subclass Peniculida (Oligohymenophorea, Ciliophora) inferred from small subunit rRNA gene sequences. *J. Eukaryot. Microbiol.* **2000**, 47, 419–429. [[CrossRef](#)]
9. Fokin, S.I.; Przyboś, E.; Chivilev, S.M.; Beier, C.L.; Horn, M.; Skotarczak, B.; Wodecka, B.; Fujishima, M. Morphological and molecular investigations of *Paramecium schewiakoffi* sp. nov. (Ciliophora, Oligohymenophorea) and current status of distribution and taxonomy of *Paramecium* spp. *Eur. J. Protistol.* **2004**, 40, 225–243. [[CrossRef](#)]
10. Pröschold, T.; Darienko, T.; Silva, P.C.; Reisser, W.; Krienitz, L. The systematics of *Zoothorella* revisited employing an integrative approach. *Environ. Microbiol.* **2011**, 13, 350–364. [[CrossRef](#)]
11. Vorobyev, K.; Andronov, E.; Rautian, M.; Skoblo, I.; Migunova, A.; Kvitko, K. An atypical *Chlorella* symbiont *Paramecium bursaria*. *Protistology* **2009**, 6, 39–44.
12. Hoshina, R.; Imamura, N. Phylogenetic position of endosymbiotic green algae in *Paramecium bursaria* Ehrenberg Japan. *Plant Biol.* **2004**, 6, 447–453. [[CrossRef](#)]
13. Gaponova, I.N.; Andronov, E.E.; Migunova, A.V.; Vorobyev, K.P.; Chizhevskaja, E.P.; Kvitko, K.V. Genomic dactyloscopy of *Chlorella* sp., symbionts of *Paramecium bursaria*. *Protistology* **2007**, 4, 311–317.
14. Migunova, A.; Kvitko, K.; Prokosheva, M.; Litvinov, D. Influence of temperature on duplication *Paramecium bursaria*, *Chlorella*, PBCV-1 viruses in system of threefold symbiosis. *Vestn. St. Peterburg Univ.* **2000**, 3, 65–75.
15. Shihira, I.; Krauss, R.W. *Chlorella: Physiology and Taxonomy of Forty-One Isolates*; University of Maryland: College Park, MD, USA, 1965; pp. 1–97.

16. Brandt, K. Ueber das Zusammenleben von Thieren und Algen. *Arch. Anat. Physiol.* **1881**, *1881*, 570–574.
17. Pröschold, T.; Bock, C.; Luo, W.; Krienitz, L. Polyphyletic distribution of bristle formation in Chlorellaceae: *Micractinium*, *Diacanthos*, *Didymogenes* and *Hegewaldia* gen. nov. (Trebouxiophyceae, Chlorophyta). *Phycol. Res.* **2010**, *58*, 1–8. [[CrossRef](#)]
18. Karakashian, S.J.; Rudzinska, M.A. Inhibition of lysosomal fusion with symbiont-containing vacuoles in *Paramecium bursaria*. *Exp. Cell Res.* **1981**, *131*, 387–393. [[CrossRef](#)]
19. Kodama, Y.; Fujishima, M. Infection of *Paramecium bursaria* by symbiotic *Chlorella* species. In *Endosymbionts in Paramecium*; Springer: Berlin, Germany, 2009; pp. 31–55.
20. Kodama, Y.; Fujishima, M. Secondary symbiosis between *Paramecium* and *Chlorella* cells. In *International Review of Cell and Molecular Biology*; Elsevier: Amsterdam, The Netherlands, 2010; Volume 279, pp. 33–77.
21. He, M.; Wang, J.; Fan, X.; Liu, X.; Shi, W.; Huang, N.; Zhao, F.; Miao, M. Genetic basis for the establishment of endosymbiosis in *Paramecium*. *ISME J.* **2019**, *13*, 1360–1369. [[CrossRef](#)]
22. Reisser, W. Die stoffwechselphysiologischen Beziehungen zwischen *Paramecium bursaria* Ehrberg und *Chlorella* spec. der *Paramecium bursaria*-Symbiose. *Arch. Microbiol.* **1976**, *107*, 357–360. [[CrossRef](#)]
23. Brown, J.A.; Nielsen, P.J. Transfer of photosynthetically produced carbohydrate from endosymbiotic *Chlorella* in *Paramecium bursaria*. *J. Protozool.* **1974**, *21*, 569–570. [[CrossRef](#)]
24. Summerer, M.; Sonntag, B.; Hörtnagl, P.; Sommaruga, R. Symbiotic ciliates receive protection against UV damage from their algae: A test with *Paramecium bursaria* and *Chlorella*. *Protist* **2009**, *160*, 233–243. [[CrossRef](#)]
25. Dunigan, D.D.; Al-Sammak, M.; Al-Ameeli, Z.; Agarkova, I.V.; DeLong, J.P.; Van Etten, J.L. Chloroviruses lure hosts through long-distance chemical signaling. *J. Virol.* **2019**, *93*, e01688-18. [[CrossRef](#)]
26. Coy, S.R.; Alsante, A.N.; Van Etten, J.L.; Wilhelm, S.W. Cryopreservation of *Paramecium bursaria* *Chlorella* Virus-1 during an active infection cycle of its host. *PLoS ONE* **2019**, *14*, e0211755. [[CrossRef](#)] [[PubMed](#)]
27. Hoshina, R.; Imamura, N. Origins of algal symbionts of *Paramecium bursaria*. In *Endosymbionts in Paramecium*; Springer: Berlin, Germany, 2009; pp. 1–29.
28. Sommaruga, R.; Sonntag, B. Photobiological aspects of the mutualistic association between *Paramecium bursaria* and *Chlorella*. In *Endosymbionts in Paramecium*; Springer: Berlin, Germany, 2009; pp. 111–130.
29. Kodama, Y.; Suzuki, H.; Dohra, H.; Sugii, M.; Kitazume, T.; Yamaguchi, K.; Shigenobu, S.; Fujishima, M. Comparison of gene expression of *Paramecium bursaria* with and without *Chlorella variabilis* symbionts. *BMC Genom.* **2014**, *15*, 183. [[CrossRef](#)] [[PubMed](#)]
30. Lowe, C.D.; Minter, E.J.; Cameron, D.D.; Brockhurst, M.A. Shining a light on exploitative host control in a photosynthetic endosymbiosis. *Curr. Biol.* **2016**, *26*, 207–211. [[CrossRef](#)] [[PubMed](#)]
31. Hoshina, R.; Fujiwara, Y. Photobiont flexibility in *Paramecium bursaria*: Double and triple photobiont co-habitation. *Adv. Microbiol.* **2012**, *2*, 227. [[CrossRef](#)]
32. Tonooka, Y.; Watanabe, T. A natural strain of *Paramecium bursaria* lacking symbiotic algae. *Eur. J. Protistol.* **2002**, *38*, 55–58. [[CrossRef](#)]
33. Nakajima, T.; Matsubara, T.; Ohta, Y.; Miyake, D. Exploitation or cooperation? Evolution of a host (ciliate)-benefiting alga in a long-term experimental microcosm culture. *Biosystems* **2013**, *113*, 127–139. [[CrossRef](#)]
34. Ishikawa, M.; Yuyama, I.; Shimizu, H.; Nozawa, M.; Ikeo, K.; Gojobori, T. Different endosymbiotic interactions in two hydra species reflect the evolutionary history of endosymbiosis. *Genome Biol. Evol.* **2016**, *8*, 2155–2163. [[CrossRef](#)]
35. Esteban, G.F.; Fenchel, T.; Finlay, B.J. Mixotrophy in ciliates. *Protist* **2010**, *161*, 621–641. [[CrossRef](#)]
36. Rautian, M.S.; Wackerow-Kouzova, N.D. Phylogenetic placement of two previously described intranuclear bacteria from the ciliate *Paramecium bursaria* (Protozoa, Ciliophora): ‘*Holospora acuminata*’ and ‘*Holospora curviuscula*’. *Int. J. Syst. Evol. Microbiol.* **2013**, *63*, 1930–1933. [[CrossRef](#)]
37. Gong, J.; Qing, Y.; Guo, X.; Warren, A. “*Candidatus Sonnebornia yantaiensis*”, a member of candidate division OD1, as intracellular bacteria of the ciliated protist *Paramecium bursaria* (Ciliophora, Oligohymenophorea). *Syst. Appl. Microbiol.* **2014**, *37*, 35–41. [[CrossRef](#)]
38. Götz, H. Infections of *Paramecium bursaria* with bacteria and yeasts. *J. Cell. Sci.* **1982**, *58*, 445–453. [[PubMed](#)]
39. Suzuki, T.; Omura, G.; Götz, H. Infection of symbiont-free *Paramecium bursaria* with yeasts. *Jpn. J. Protozool.* **2003**, *36*, 17–18.

40. Nakahara, M.; Handa, S.; Watanabe, S.; Deguchil, H. *Choricystis minor* as a new symbiont of simultaneous two-species association with *Paramecium bursaria* and implications for its phylogeny. *Symbiosis* **2004**, *36*, 127–151.
41. Szokoli, F.; Sabaneyeva, E.; Castelli, M.; Krenek, S.; Schrallhammer, M.; Soares, C.A.; da Silva-Neto, I.D.; Berendonk, T.U.; Petroni, G. “*Candidatus Fokinia solitaria*”, a novel “stand-alone” symbiotic lineage of *Midichloriaceae* (*Rickettsiales*). *PLoS ONE* **2016**, *11*, e0145743. [[CrossRef](#)]
42. Krenek, S.; Berendonk, T.U.; Fokin, S.I. New *Paramecium* (Ciliophora, Oligohymenophorea) congeners shape our view of its biodiversity. *Org. Divers. Ecol.* **2015**, *15*, 215–233. [[CrossRef](#)]
43. Da Silva Paiva, T.; do Nascimento Borges, B.; Harada, M.L.; da Silva-Neto, I.D. Description and molecular phylogeny of *Paramecium grohmannae* sp. nov. (Ciliophora, Peniculida) from a wastewater treatment plant in Brazil. *Rev. Bras. De Zoociências* **2016**, *17*, 7–19.
44. Castelli, M.; Serra, V.; Senra, M.V.; Basuri, C.K.; Soares, C.A.; Fokin, S.I.; Modeo, L.; Petroni, G. The hidden world of *Rickettsiales* symbionts: “*Candidatus Spectririckettsia obscura*,” a novel bacterium found in Brazilian and Indian *Paramecium caudatum*. *Microb. Ecol.* **2019**, *77*, 748–758. [[CrossRef](#)]
45. Sonneborn, T.M. Mating types in *Paramecium aurelia*: Diverse conditions for mating in different stocks; occurrence, number and interrelations of the types. *Proc. Am. Philosoph. Soc.* **1938**, *79*, 411–434.
46. Bomford, R. The syngens of *Paramecium bursaria*: New mating types and intersyngenic mating reactions. *J. Protozool.* **1966**, *13*, 497–501. [[CrossRef](#)]
47. Greczek-Stachura, M.; Potekhin, A.; Przyboś, E.; Rautian, M.; Skoblo, I.; Tarcz, S. Identification of *Paramecium Bursaria* syngens through molecular markers—comparative analysis of three loci in the nuclear and mitochondrial DNA. *Protist* **2012**, *163*, 671–685. [[CrossRef](#)]
48. Strüder-Kypke, M.C.; Lynn, D.H. Comparative analysis of the mitochondrial cytochrome c oxidase subunit I (COI) gene in ciliates (Alveolata, Ciliophora) and evaluation of its suitability as a biodiversity marker. *Syst. Biodivers.* **2010**, *8*, 131–148. [[CrossRef](#)]
49. Tasneem, F.; Shakoori, F.R.; Ilyas, M.; Shahzad, N.; Potekhin, A.; Shakoori, A.R. Genetic diversity of *Paramecium* species on the basis of multiple loci analysis and ITS secondary structure models. *J. Cell. Biochem.* **2019**. [[CrossRef](#)] [[PubMed](#)]
50. Spanner, C.; Darienko, T.; Biebler, T.; Sonntag, B.; Pröschold, T. Endosymbiotic green algae in *Paramecium Bursaria*: A new isolation method and a simple diagnostic PCR approach for the identification. *Diversity* **2020**, *12*, 240. [[CrossRef](#)]
51. Potekhin, A.; Mayén-Estrada, R. *Paramecium* diversity and a new member of the *Paramecium Aurelia* species complex described from Mexico. *Diversity* **2020**, *12*, 197. [[CrossRef](#)]
52. Hoshina, R.; Fujiwara, Y. Molecular characterization of *Chlorella* cultures of the National Institute for Environmental Studies Culture Collection with description of *Micractinium inernum* sp. nov., *Didymogonus sphaerica* sp. nov., and *Didymogonus soliella* sp. nov. (Chlorellaceae, Trebouxiophyceae). *Phycol. Res.* **2013**, *61*, 124–132.
53. Krienitz, L.; Bock, C. Present state of the systematics of planktonic coccoid green algae of inland waters. *Hydrobiologia* **2012**, *698*, 295–326. [[CrossRef](#)]
54. Luo, W.; Pröschold, T.; Bock, C.; Krienitz, L. Generic concept in *Chlorella*-related coccoid green algae (Chlorophyta, Trebouxiophyceae). *Plant Biol.* **2010**, *12*, 545–553. [[CrossRef](#)]
55. Pröschold, T.; Pitsch, G.; Darienko, T. *Micractinium tetrahymenae* (Trebouxiophyceae, Chlorophyta), a new endosymbiont isolated from ciliates. *Diversity* **2020**, *12*, 200. [[CrossRef](#)]
56. Heeg, J.S.; Wolf, M. ITS2 and 18S rDNA sequence-structure phylogeny of *Chlorella* and allies (Chlorophyta, Trebouxiophyceae, Chlorellaceae). *Plant Gene* **2015**, *4*, 20–28. [[CrossRef](#)]
57. Krienitz, L.; Hegewald, E.H.; Hepperle, D.; Huss, V.A.; Rohr, T.; Wolf, M. Phylogenetic relationship of *Chlorella* and *Parachlorella* gen. nov. (Chlorophyta, Trebouxiophyceae). *Phycology* **2004**, *43*, 529–542. [[CrossRef](#)]
58. Krienitz, L.; Bock, C.; Kotut, K.; Pröschold, T. Genotypic diversity of *Dictyosphaerium*-morphospecies (Chlorellaceae, Trebouxiophyceae) in African inland waters, including the description of four new genera. *Fottea* **2012**, *12*, 231–253. [[CrossRef](#)]
59. Bella, C.; Koehler, L.; Grosser, K.; Berendonk, T.U.; Petroni, G.; Schrallhammer, M. Fitness impact of obligate intranuclear bacterial symbionts depends on host growth phase. *Front. Microbiol.* **2016**, *7*, 2084. [[CrossRef](#)] [[PubMed](#)]

60. Zagata, P.; Greczek-Stachura, M.; Tarcz, S.; Rautian, M. The evolutionary relationships between endosymbiotic green algae of *Paramecium bursaria* syngens originating from different geographical locations. *Folia Biol.* **2016**, *64*, 47–54. [CrossRef] [PubMed]
61. Fokin, S.; Andreoli, I.; Verni, F.; Petroni, G. *Apofrontonia dohrni* sp. n. and the phylogenetic relationships within Peniculia (Protista, Ciliophora, Oligohymenophorea). *Zool. Scr.* **2006**, *35*, 289–300. [CrossRef]
62. Darienko, T.; Rad-Menéndez, C.; Campbell, C.; Pröschold, T. Are there any true marine *Chlorella* species? Molecular phylogenetic assessment and ecology of marine *Chlorella*-like organisms, including a description of *Droopiella* gen. nov. *Syst. Biodivers.* **2019**, *17*, 811–829. [CrossRef] [PubMed]
63. Marin, M.; Castro, B.; Gaitan, A.; Preisig, O.; Wingfield, B.; Wingfield, M. Relationships of *Ceratocystis fimbriata* isolates from Colombian coffee-growing regions based on molecular data and pathogenicity. *J. Phytopathol.* **2003**, *151*, 395–405. [CrossRef]
64. Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F.O. The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* **2012**, *41*, D590–D596. [CrossRef]
65. Swofford, D. *PAUP**. *Phylogenetic analysis using parsimony (* and other, odes)*. Version 4.0 b10 for Macintosh; Sinauer Associates Inc.: Sunderland, MA, USA, 2002.
66. Huelsenbeck, J.P.; Ronquist, F. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **2001**, *17*, 754–755. [CrossRef]
67. Guindon, S.; Gascuel, O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **2003**, *52*, 696–704. [CrossRef]
68. Darienko, T.; Gustavs, L.; Pröschold, T. Species concept and nomenclatural changes within the genera *Elliptochloris* and *Pseudochlorella* (Trebouxiophyceae) based on an integrative approach. *J. Phycol.* **2016**, *52*, 1125–1145. [CrossRef]
69. Chae, H.; Lim, S.; Kim, H.S.; Choi, H.G.; Kim, J.H.; Chae, H.; Lim, S.; Kim, H.S.; Choi, H.G.; Kim, J.H. Morphology and phylogenetic relationships of *Micractinium* (Chlorellaceae, Trebouxiophyceae) taxa, including three new species from Antarctica. *Algae* **2019**, *34*, 267–275. [CrossRef]
70. Hodač, L.; Hallmann, C.; Spitzer, K.; Elster, J.; Faßhauer, F.; Brinkmann, N.; Lepka, D.; Diwan, V.; Friedl, T. Widespread green algae *Chlorella* and *Stichococcus* exhibit polar-temperate and tropical-temperate biogeography. *FEMS Microbiol. Ecol.* **2016**, *92*, 1–16. [CrossRef] [PubMed]
71. Mathews, D.H.; Burkard, M.E.; Freier, S.M.; Wyatt, J.R.; Turner, D.H. Predicting oligonucleotide affinity to nucleic acid targets. *RNA* **1999**, *5*, 1458–1469. [CrossRef] [PubMed]
72. Meier, R.; Wiessner, W. Infection of algae-free *Paramecium bursaria* symbiotic *Chlorella* sp. isolated from green paramecia: II. A timed study. *J. Cell Sci.* **1989**, *93*, 571–579. [CrossRef]
73. Kodama, Y.; Fujishima, M. Characteristics of the digestive vacuole membrane of the alga-bearing ciliate *Paramecium bursaria*. *Protist* **2012**, *163*, 658–670. [CrossRef]
74. Hoshina, R.; Kato, Y.; Kamako, S.; Imamura, N. Genetic evidence of “American” and “European” type symbiotic algae of *Paramecium bursaria* Ehrenberg. *Plant Biol.* **2005**, *7*, 526–532. [CrossRef]
75. Hoshina, R.; Imamura, N. Multiple origins of the symbioses in *Paramecium bursaria*. *Protist* **2008**, *159*, 53–63. [CrossRef]
76. Hoshina, R.; Iwataki, M.; Imamura, N. *Chlorella variabilis* and *Micractinium reisseri* sp. nov. (Chlorellaceae, Trebouxiophyceae): Redescription of the endosymbiotic green algae of *Paramecium bursaria* (Peniculia, Oligohymenophorea) in the 120th year. *Phycol. Res.* **2010**, *58*, 188–201. [CrossRef]
77. Summerer, M.; Sonntag, B.; Sommaruga, R. An experimental test of the symbiosis specificity between the ciliate *Paramecium bursaria* and strains of the unicellular green alga *Chlorella*. *Environ. Microbiol.* **2007**, *9*, 2117–2122. [CrossRef]

