

“Zakopane” AMPERE NMR SCHOOL
virtual event

BOOK OF ABSTRACTS

Edited by S. Jurga
Adam Mickiewicz University in Poznań
NanoBioMedical Centre

21st June – 23rd July 2021

“Zakopane” AMPERE NMR SCHOOL

virtual event

21st June – 23rd July 2021

organized by

**NanoBioMedical Centre
and The Centre for European Integration
Adam Mickiewicz University in Poznań, Poznań**




Organizing Committee

S. Jurga – Chair
L. Szutkowska – Executive Secretary
D. Flak
P. Florczak
B. Grześkowiak
M. Jarek
M. Jancelewicz
J. Jencyk
B. Kawczyński
G. Nowaczyk
R. Markiewicz
K. Szutkowski
A. Woźniak
T. Zalewski
K. Załęski

Scientific Committee

B. Blümich (Aachen), Germany
J. Dolinšek (Ljubljana), Slovenia
S. Jurga (Poznań), Poland
W. Koźmiński (Warszawa), Poland
D. Kruk (Olsztyn) Poland
D. Lurie (Aberdeen), UK
A. MacKay (Vancouver), Canada
B. Meier (Zürich), Switzerland

**THIS SCHOOL IS SUPPORTED BY
GROUPEMENT AMPERE, SOFTCOMP, EUSMI**



SPEAKERS	7
MOLECULAR DYNAMICS BY NMR FOR MATERIALS TESTING: RELAXATION, EXCHANGE AND MRI <u>Bernhard Blümich</u>	9
BASICS OF MRI AND RESEARCH ON FAST FIELD-CYCLING MRI <u>David J. Lurie, Lionel M. Broche, Gareth R. Davies, Mary J. MacLeod, P. James Ross, Robert Stormont</u>	10
THE HIGHER, THE BETTER?! THE PROMISES AND PERKS OF ULTRA-HIGH FIELD MRI AT 22.3 T <u>Julia R. Krug</u>	11
DIFFUSION-RELAXATION CORRELATION MRI <u>D. Topgaard</u>	12
NMR STUDIES OF COMPLEX SAMPLES USING COMPACT INSTRUMENTS <u>Anton Duchowny</u>	13
INSIGHT INTO THE DETAILS OF MOLECULAR TRANSLATION DYNAMICS IN LIQUIDS BY NMR GRADIENT SPIN ECHO METHOD <u>Janez Stepišnik, Aleš Mohorič</u>	14
ULTRAFast MULTIDIMENSIONAL RELAXATION AND DIFFUSION MEASUREMENTS <u>Ville-Veikko Telkki</u>	15
MEASURING NMR RELAXATION TIMES – WHAT CAN POSSIBLY GO WRONG!? <u>Siegfried Stapf</u>	16
WHY ARE SOME NUCLEI NON-SPHERICAL AND WHAT DOES THAT HAVE TO DO WITH SPIN? <u>Ilya Kuprov</u>	17
RESIDUAL LINE WIDTH IN MAS SOLID-STATE NMR <u>Matthias Ernst</u>	18
SOLID-STATE NMR OF QUADRUPOLEAR NUCLEI AND THEIR NEIGHBORS <u>Olivier Lafon</u>	19
HIGH-ENTROPY ALLOYS <u>Janez Dolinšek</u>	20
THE KINETICS OF PROTON TRANSPORT AND THERMAL PROCESSES IN ANHYDROUS NANOCOMPOSITE PROTON CONDUCTOR BASED ON CELLULOSE <u>Michał Bielejewski</u>	21
NMR STUDIES OF POLYMER GEL ELECTROLYTES <u>Claudia Schmidt, Waldemar Keil, Martin Siebrecht, Shilpa Khurana, Manoj K. Vyas, Amita Chandra</u>	22
CONFORMATIONAL AND AGGREGATIONAL BEHAVIOR OF SOME SURFACTANTS IN AQUEOUS SOLUTIONS BY NMR OF ¹H AND ¹³C NUCLEI <u>V. I. Chizhik, M. V. Popova, A. V. Ievlev, V. V. Vasinovich</u>	23
CARBON- AND PROTON-DETECTED SOLID-STATE NMR SEQUENTIAL ASSIGNMENTS AND APPLICATIONS TO FIBRILS AND MEMBRANE PROTEINS <u>Anja Böckmann</u>	24
FAST MAS AND BIOMOLECULES <u>Beat H. Meier, Morgane Callon, Alexander A. Malär, Gyula Palfy, Sara Pfister, Václav Rimal, Marco E. Weber, Thomas Wiegand, Johannes Zehnder, Matias Chávez, Lauriane Lecoq, Matthias Ernst, Anja Böckman, Beat H. Meier</u>	25
NMR STUDIES OF SOLID NANOCRYSTALLINE CELLULOSE-BASED PROTON CONDUCTORS <u>Jadwiga Tritt-Goc</u>	26
HIGH DIMENSIONALITY AND HIGH RESOLUTION NMR EXPERIMENTS FOR BIOMOLECULES <u>Wiktor Koźmiński</u>	27
EXPLORING SCALAR COUPLINGS AND CHEMICAL EXCHANGE FROM LOW TO ULTRA-HIGH FIELDS <u>Fabien Ferrage</u>	28

CONFORMATIONAL AND AGGREGATIONAL BEHAVIOR OF SOME SURFACTANTS IN AQUEOUS SOLUTIONS BY NMR OF ^1H AND ^{13}C NUCLEI

V. I. Chizhik, M. V. Popova, A. V. Ievlev, and V. V. Vasinovich

*Saint-Petersburg State University, Faculty of Physics, 198504 Saint Petersburg,
Uliyanovskaya st., 1, v.chizhik@spbu.ru*

Surfactants belong to actual chemical products, which are consumed in a large amount and find numerous applications in various areas. Amino-acid based surfactants attract an increasing interest, mainly because of their environmentally friendly properties. In comparison to conventional surfactants they are characterized in particular by a fast biodegradation and excellent antimicrobial and antifungal activities. Among them one of the most commonly used is sodium N-lauroyl sarcosinate (SLS).

To investigate the chemical and physical properties of surfactant solutions, a great number of experimental and theoretical methods have been used. Among them a special place can be assigned to nuclear magnetic resonance methods (used in the present study): NMR spectroscopy, NMR relaxometry, NMR diffusometry. Despite numerous studies of the properties of SLS in various systems, there are currently many questions that require clarification, in particular it concerns the value of the critical micelle concentration (CMC), conformational transitions in solutions and mixtures, local mobility of individual segments of amphiphilic molecules, etc. For example, as reported in the literature, the value of CMC for SLS lies in the range between 0.009 and 0.014 M, depending on the used method.

From the spectra it was possible to estimate the time scale of the internal reorientations of the COO-group at the room temperature: 3–5 ms. There is a relatively fast exchange of the SLS molecules between the monomer and micellar states: the exchange time is much less than 1 ms. Modern NMR spectrometers allow the measurements of relaxation rates for resolved spectral lines belonging to different molecular groups and, therefore, the investigation the local molecular mobility. For example, it was possible to obtain for the set of CH_2 groups the correlation time of $\tau_c \sim 10^{-10}$ s. The bigger values of spin-relaxation times for the CH_3 groups indicate a higher mobility of these groups. Using the NMR method with a pulsed magnetic field gradient, the self-diffusion coefficients of surfactant molecules in the monomeric and micellar states were measured. The obtained data allowed us to refine the CMC values and develop a new method for estimating the decrease in the concentration of monomers with an increase in the total concentration of surfactants.

In practical applications the surfactant mixtures are often used because surfactants in mixtures exhibit new properties. In particular, mixtures of cationic and anionic surfactants are of interest for research. In the present study, in addition to SLS the dodecyltrimethylammonium bromide (DTAB) was chosen as the starting point. Data indicating the formation of mixed micelles in SLS+DTAB systems have been obtained. As a hypothesis, we suggest the electrostatic binding of molecules in the SLS+DTAB systems, which manifests itself in the broadening of the spectral lines for certain concentration regions. The spin-lattice relaxation times of ^1H and ^{13}C nuclei have been measured. Based on the relaxation of ^{13}C , the reorientation times of the molecules are calculated. The bigger correlation times were observed for some molecular groups that correlate with the broadening of lines in the ^1H spectrum for these groups.