



XXVIII  
МЕЖДУНАРОДНАЯ  
ЧУГАЕВСКАЯ КОНФЕРЕНЦИЯ  
ПО КООРДИНАЦИОННОЙ ХИМИИ

# СБОРНИК ТЕЗИСОВ

*XXVIII Международная Чугаевская конференция  
по Координационной химии*

*XVIII Международная конференция  
«Спектроскопия координационных соединений»*

*V Молодежная школа-конференция  
«Физико-химические методы в химии координаци-  
онных соединений»*

03 - 08 октября 2021 года,  
Туапсе, Ольгинка, Краснодарский край, Россия

---

по Координационной химии

V Молодежная школа-конференция  
«Физико-химические методы в химии координационных соединений»

ПЛЕНАРНЫЕ ДОКЛАДЫ .....	10
УСТНЫЕ ДОКЛАДЫ	
XXVIII Международная Чугаевская конференция по Координационной химии .....	37
V Молодежная школа-конференция «Физико-химические методы в химии координационных соединений» .....	152
ПОСТЕРНЫЕ ДОКЛАДЫ .....	225
ЗАОЧНЫЕ ДОКЛАДЫ .....	329

Сборник тезисов

XVIII Международной конференции  
«Спектроскопия координационных соединений»

МЕТОДЫ ИССЛЕДОВАНИЯ И ПОЛУЧЕНИЯ СУПРА-И НАНОСИСТЕМ .....	468
МЕТОДЫ ОПТИЧЕСКОЙ СПЕКТРОСКОПИИ И КВАНТОВОЙ ХИМИИ В ИССЛЕДОВАНИИ КС .....	480
СИНТЕЗ, СПЕКТРАЛЬНЫЕ СВОЙСТВА И ПРИМЕНЕНИЕ КОМПЛЕКСНЫХ СОЕДИНЕНИЙ .....	494
ФОТОХИМИЯ И ЛЮМИНЕСЦЕНЦИЯ КООРДИНАЦИОННЫХ СОЕДИНЕНИЙ .....	518
ЯМР И ЭПР СПЕКТРОСКОПИЯ КООРДИНАЦИОННЫХ СОЕДИНЕНИЙ .....	541



# СБОРНИК ТЕЗИСОВ

Сборник тезисов  
XVIII Международной конференции  
«Спектроскопия координационных соединений»

Туапсе,  
3-8 октября 2021 года



**ЯМР И ЭПР СПЕКТРОСКОПИЯ  
КООРДИНАЦИОННЫХ  
СОЕДИНЕНИЙ**



## CONFORMATIONAL AND AGGREGATIONAL BEHAVIOR OF SODIUM N-LAUROYL SARCOSINATE IN AQUEOUS SOLUTIONS BY NMR OF $^1\text{H}$ AND $^{13}\text{C}$ NUCLEI

V.I. Chizhik<sup>a</sup>, M.V. Popova<sup>b</sup>, V.V. Vasinovich<sup>a</sup>, and A.V. Ievlev<sup>a</sup>

<sup>a</sup>*Saint-Petersburg State University, Saint Petersburg, Russian Federation*  
<sup>b</sup>*Konstatinov Institute of Nuclear Physics of the NRC "Kurchatov Institute",  
Gatchina, 188300 Russian Federation*

Surfactants belong to actual chemical products, which are consumed in a large amount and find numerous applications in various areas. Amino-acid based surfactants attract an increasing interest, mainly because of their environmentally friendly properties. Among them one of the most commonly used is sodium N-lauroyl sarcosinate (SLS).

To investigate the chemical and physical properties of surfactant solutions, a great number of experimental and theoretical methods have been used. Among them a special place can be assigned to nuclear magnetic resonance methods (used in the present study): NMR spectroscopy, NMR relaxometry, NMR diffusometry. Despite numerous studies of the properties of SLS in various systems, there are currently many questions that require clarification, in particular it concerns the value of the critical micelle concentration (CMC), conformational transitions in solutions and mixtures, local mobility of individual segments of amphiphilic molecules, etc.

From the NMR spectra it was possible to estimate the time scale of the internal reorientations of the COO-group at the room temperature: 3÷5 ms. There is a relatively fast exchange of the SLS molecules between the monomer and micellar states: the exchange time is much less than 1 ms. Modern NMR spectrometers allow the measurements of relaxation rates for resolved spectral lines belonging to different molecular groups and, therefore, the investigation the local molecular mobility. For example, it was possible to obtain for the set of  $\text{CH}_2$  groups the correlation time of  $\tau_c \sim 10^{-10}$  s. Using the NMR method with a pulsed magnetic field gradient, the self-diffusion coefficients of surfactant molecules in the monomeric and micellar states were measured. The obtained data allowed us to refine the CMC values and develop a new method for estimating the decrease in the concentration of monomers with the increase in the total concentration of surfactants.

In practical applications the surfactant mixtures are often used because surfactants in mixtures exhibit new properties. In particular, mixtures of cationic and anionic surfactants are of interest for research. In the present study, in addition to SLS the dodecyltrimethylammonium bromide (DTAB) was chosen as the starting point. Data indicating the formation of mixed micelles in SLS+DTAB systems have been obtained. As a hypothesis, we suggest the electrostatic binding of molecules in the SLS+DTAB systems, which manifests itself in the broadening of the spectral lines for certain concentration regions. The spin-lattice relaxation times of  $^1\text{H}$  and  $^{13}\text{C}$  nuclei have been measured. Based on the relaxation of  $^{13}\text{C}$ , the reorientation times of the molecules are calculated.

**Acknowledgements:** All NMR measurements were performed at the Research park of St. Petersburg State University, «Center for Magnetic Resonance».

**e-mail:** v.chizhik@spbu.ru